

Leavitt path algebras: Something for everyone

Gene Abrams



AMSI AGR National Seminar Series

February 8, 2013

Overview

- 1 Leavitt path algebras: Introduction and Motivation
- 2 Algebraic properties
- 3 Projective modules
- 4 Connections and Applications

- 1 Leavitt path algebras: Introduction and Motivation
- 2 Algebraic properties
- 3 Projective modules
- 4 Connections and Applications

Brief history, and motivating examples

One of the first theorems you saw as an undergraduate student:

Dimension Theorem for Vector Spaces. Every nonzero vector space V has a basis. Moreover, if \mathcal{B} and \mathcal{B}' are two bases for V , then $|\mathcal{B}| = |\mathcal{B}'|$.

Brief history, and motivating examples

One of the first theorems you saw as an undergraduate student:

Dimension Theorem for Vector Spaces. Every nonzero vector space V has a basis. Moreover, if \mathcal{B} and \mathcal{B}' are two bases for V , then $|\mathcal{B}| = |\mathcal{B}'|$.

Note: V has a basis $\mathcal{B} = \{b_1, b_2, \dots, b_n\} \Leftrightarrow V \cong \bigoplus_{i=1}^n \mathbb{R}$ as vector spaces. So:

One result of Dimension Theorem, Rephrased:

$$\bigoplus_{i=1}^n \mathbb{R} \cong \bigoplus_{i=1}^m \mathbb{R} \Leftrightarrow m = n.$$

Brief history, and motivating examples

The same Dimension Theorem holds, with the identical proof, if K is any division ring (i.e., any ring for which every nonzero element has a multiplicative inverse).

Brief history, and motivating examples

The same Dimension Theorem holds, with the identical proof, if K is any division ring (i.e., any ring for which every nonzero element has a multiplicative inverse).

Idea: Show any maximal linearly independent subset of V actually spans V . Why are there *any* linearly independent subsets at all?

Brief history, and motivating examples

The same Dimension Theorem holds, with the identical proof, if K is any division ring (i.e., any ring for which every nonzero element has a multiplicative inverse).

Idea: Show any maximal linearly independent subset of V actually spans V . Why are there *any* linearly independent subsets at all?

If $0 \neq v \in V$, then $\{v\}$ is linearly independent.

Brief history, and motivating examples

The same Dimension Theorem holds, with the identical proof, if K is any division ring (i.e., any ring for which every nonzero element has a multiplicative inverse).

Idea: Show any maximal linearly independent subset of V actually spans V . Why are there *any* linearly independent subsets at all?

If $0 \neq v \in V$, then $\{v\}$ is linearly independent.

If $kv = 0$, need to show $k = 0$. But $k \neq 0 \Rightarrow \frac{1}{k}kv = 0 \Rightarrow v = 0$, contradiction.

Similar idea (multiply by the inverse of a nonzero element of K) shows that a maximal linearly independent subset of V actually spans V .

Brief history, and motivating examples

Rings: e.g., \mathbb{Z} , $M_2(\mathbb{R})$, $C(\mathbb{R})$, ...

Brief history, and motivating examples

Rings: e.g., \mathbb{Z} , $M_2(\mathbb{R})$, $C(\mathbb{R})$, ...

Vector spaces for rings = “modules”

Brief history, and motivating examples

Rings: e.g., \mathbb{Z} , $M_2(\mathbb{R})$, $C(\mathbb{R})$, ...

Vector spaces for rings = “modules”

Brief history, and motivating examples

Rings: e.g., \mathbb{Z} , $M_2(\mathbb{R})$, $C(\mathbb{R})$, ...

Vector spaces for rings = “modules”

For R any ring, $n \in \mathbb{N}$, then $\bigoplus_{i=1}^n R$ is an R -module as usual.

$$r \cdot (r_1, r_2, \dots, r_n) = (rr_1, rr_2, \dots, rr_n)$$

Brief history, and motivating examples

Rings: e.g., \mathbb{Z} , $M_2(\mathbb{R})$, $C(\mathbb{R})$, ...

Vector spaces for rings = “modules”

For R any ring, $n \in \mathbb{N}$, then $\bigoplus_{i=1}^n R$ is an R -module as usual.

$$r \cdot (r_1, r_2, \dots, r_n) = (rr_1, rr_2, \dots, rr_n)$$

For a module P over a ring R we can still talk about a *basis* for P . (Note: in general, not all modules *have* bases; those that do are called *free* R -modules.)

$\bigoplus_{i=1}^n R$ always has a basis having n elements, e.g.,

$$\{e_1 = (1, 0, 0, \dots, 0), e_2 = (0, 1, 0, \dots, 0), \dots, e_n = (0, 0, 0, \dots, 1)\}$$

Brief history, and motivating examples

Question: Is the Dimension Theorem true for rings in general?

That is, if R is a ring, and $\bigoplus_{i=1}^n R \cong \bigoplus_{i=1}^m R$ as R -modules, must $m = n$?

Brief history, and motivating examples

Question: Is the Dimension Theorem true for rings in general? That is, if R is a ring, and $\bigoplus_{i=1}^n R \cong \bigoplus_{i=1}^m R$ as R -modules, must $m = n$?

Answer: NO

(But the answer is YES for the rings \mathbb{Z} , $M_2(\mathbb{R})$, $C(\mathbb{R})$)

Example: Consider the ring S of linear transformations from an infinite dimensional \mathbb{R} -vector space V to itself.

Brief history, and motivating examples

Question: Is the Dimension Theorem true for rings in general? That is, if R is a ring, and $\bigoplus_{i=1}^n R \cong \bigoplus_{i=1}^m R$ as R -modules, must $m = n$?

Answer: NO

(But the answer is YES for the rings \mathbb{Z} , $M_2(\mathbb{R})$, $C(\mathbb{R})$)

Example: Consider the ring S of linear transformations from an infinite dimensional \mathbb{R} -vector space V to itself.

Think of V as $\bigoplus_{i=1}^{\infty} \mathbb{R}$. Then think of S as $\text{RFM}(\mathbb{R})$.

Brief history, and motivating examples

Intuitively, S and $S \oplus S$ have a chance to be “the same”.

$M \mapsto$ (Odd numbered columns of M , Even numbered columns of M)

Brief history, and motivating examples

Intuitively, S and $S \oplus S$ have a chance to be “the same”.

$M \mapsto$ (Odd numbered columns of M , Even numbered columns of M)

More formally: Let

$$Y_1 = \begin{pmatrix} 1 & 0 & 0 & 0 & 0 & 0 & \dots \\ 0 & 0 & 0 & 0 & 0 & 0 & \dots \\ 0 & 1 & 0 & 0 & 0 & 0 & \dots \\ 0 & 0 & 0 & 0 & 0 & 0 & \dots \\ 0 & 0 & 1 & 0 & 0 & 0 & \dots \\ 0 & 0 & 0 & 0 & 0 & 0 & \dots \\ \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \ddots \end{pmatrix} \quad Y_2 = \begin{pmatrix} 0 & 0 & 0 & 0 & 0 & 0 & \dots \\ 1 & 0 & 0 & 0 & 0 & 0 & \dots \\ 0 & 0 & 0 & 0 & 0 & 0 & \dots \\ 0 & 1 & 0 & 0 & 0 & 0 & \dots \\ 0 & 0 & 0 & 0 & 0 & 0 & \dots \\ 0 & 0 & 1 & 0 & 0 & 0 & \dots \\ \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \ddots \end{pmatrix}$$

$$X_1 = \begin{pmatrix} 1 & 0 & 0 & 0 & 0 & 0 & \dots \\ 0 & 0 & 1 & 0 & 0 & 0 & \dots \\ 0 & 0 & 0 & 0 & 1 & 0 & \dots \\ 0 & 0 & 0 & 0 & 0 & 0 & \dots \\ 0 & 0 & 0 & 0 & 0 & 0 & \dots \\ 0 & 0 & 0 & 0 & 0 & 0 & \dots \\ \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \ddots \end{pmatrix} \quad X_2 = \begin{pmatrix} 0 & 1 & 0 & 0 & 0 & 0 & \dots \\ 0 & 0 & 0 & 1 & 0 & 0 & \dots \\ 0 & 0 & 0 & 0 & 0 & 1 & \dots \\ 0 & 0 & 0 & 0 & 0 & 0 & \dots \\ 0 & 0 & 0 & 0 & 0 & 0 & \dots \\ 0 & 0 & 0 & 0 & 0 & 0 & \dots \\ \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \ddots \end{pmatrix}$$

Brief history, and motivating examples

Then MY_1 gives the Odd Columns of M , while MY_2 gives the Even Columns of M .

Brief history, and motivating examples

Then MY_1 gives the Odd Columns of M , while MY_2 gives the Even Columns of M .

So the previous intuitive map is, formally, $M \mapsto (MY_1, MY_2)$.

Brief history, and motivating examples

Then MY_1 gives the Odd Columns of M , while MY_2 gives the Even Columns of M .

So the previous intuitive map is, formally, $M \mapsto (MY_1, MY_2)$.

Similarly, we should be able to 'go back' from pairs of matrices to a single matrix, by interweaving the columns.

Brief history, and motivating examples

Then MY_1 gives the Odd Columns of M , while MY_2 gives the Even Columns of M .

So the previous intuitive map is, formally, $M \mapsto (MY_1, MY_2)$.

Similarly, we should be able to 'go back' from pairs of matrices to a single matrix, by interweaving the columns.

That is, more formally, $(M_1, M_2) \mapsto M_1X_1 + M_2X_2$ is a reasonable way to associate a pair of matrices with a single one.

Brief history, and motivating examples

Here's what's really going on. These equations are easy to verify:

$$Y_1 X_1 + Y_2 X_2 = I,$$

$$X_1 Y_1 = I = X_2 Y_2, \quad \text{and} \quad X_1 Y_2 = 0 = X_2 Y_1.$$

Brief history, and motivating examples

Here's what's really going on. These equations are easy to verify:

$$Y_1 X_1 + Y_2 X_2 = I,$$

$$X_1 Y_1 = I = X_2 Y_2, \quad \text{and} \quad X_1 Y_2 = 0 = X_2 Y_1.$$

Using these, we get inverse maps:

$$S \rightarrow S \oplus S \quad \text{via} \quad M \mapsto (MY_1, MY_2), \quad \text{and}$$

$$S \oplus S \rightarrow S \quad \text{via} \quad (M_1, M_2) \mapsto M_1 X_1 + M_2 X_2.$$

For example:

$$M \mapsto (MY_1, MY_2) \mapsto MY_1 X_1 + MY_2 X_2 = M \cdot I = M.$$

Brief history, and motivating examples

Using exactly the same idea, let R be ANY ring which contains four elements y_1, y_2, x_1, x_2 satisfying

$$y_1x_1 + y_2x_2 = 1_R,$$

$$x_1y_1 = 1_R = x_2y_2, \quad \text{and} \quad x_1y_2 = 0 = x_2y_1.$$

Then $R \cong R \oplus R$.

Brief history, and motivating examples

Remark: Here the sets $\{1_R\}$ and $\{x_1, x_2\}$ are each bases for R .

Brief history, and motivating examples

Remark: Here the sets $\{1_R\}$ and $\{x_1, x_2\}$ are each bases for R .

Actually, when $R \cong R \oplus R$ as R -modules, then $\bigoplus_{i=1}^m R \cong \bigoplus_{i=1}^n R$ for *all* $m, n \in \mathbb{N}$.

Leavitt algebras

Natural question:

Does there exist R with, e.g., $R \cong R \oplus R \oplus R$, but $R \not\cong R \oplus R$?

Leavitt algebras

Natural question:

Does there exist R with, e.g., $R \cong R \oplus R \oplus R$, but $R \not\cong R \oplus R$?

Theorem

(William G. Leavitt, Trans. Amer. Math. Soc., 1962)

For every $m < n \in \mathbb{N}$ and field K there exists a K -algebra $R = L_K(m, n)$ with $\bigoplus_{i=1}^m R \cong \bigoplus_{i=1}^n R$, and all isomorphisms between free left R -modules result precisely from this one. Moreover, $L_K(m, n)$ is universal with this property.

Leavitt algebras

The $m = 1$ situation of Leavitt's Theorem is now somewhat familiar. Similar to the $n = 2$ case that we saw above, $R \cong R^n$ if and only if there exist

$$x_1, x_2, \dots, x_n, y_1, y_2, \dots, y_n \in R$$

for which

$$\sum_{i=1}^n y_i x_i = 1_R \quad \text{and} \quad x_i y_j = \delta_{i,j} 1_R.$$

Leavitt algebras

The $m = 1$ situation of Leavitt's Theorem is now somewhat familiar. Similar to the $n = 2$ case that we saw above, $R \cong R^n$ if and only if there exist

$$x_1, x_2, \dots, x_n, y_1, y_2, \dots, y_n \in R$$

for which

$$\sum_{i=1}^n y_i x_i = 1_R \quad \text{and} \quad x_i y_j = \delta_{i,j} 1_R.$$

$L_K(1, n)$ is the quotient

$$K \langle X_1, X_2, \dots, X_n, Y_1, Y_2, \dots, Y_n \rangle / \langle \left(\sum_{i=1}^n Y_i X_i \right) - 1_K; X_i Y_j - \delta_{i,j} 1_K \rangle$$

Note: $\text{RFM}(K)$ is much bigger than $L_K(1, 2)$.



Leavitt algebras

As a result, we have: Let S denote $L_K(1, n)$. Then

$$S^a \cong S^b \Leftrightarrow a \equiv b \pmod{n-1}.$$

In particular, $S \cong S^n$.

Leavitt algebras

As a result, we have: Let S denote $L_K(1, n)$. Then

$$S^a \cong S^b \Leftrightarrow a \equiv b \pmod{n-1}.$$

In particular, $S \cong S^n$.

It turns out:

Theorem. (Leavitt, Duke J. Math, 1964)

For every field K and $n \geq 2$, $L_K(1, n)$ is simple.

Leavitt algebras

As a result, we have: Let S denote $L_K(1, n)$. Then

$$S^a \cong S^b \Leftrightarrow a \equiv b \pmod{n-1}.$$

In particular, $S \cong S^n$.

It turns out:

Theorem. (Leavitt, Duke J. Math, 1964)

For every field K and $n \geq 2$, $L_K(1, n)$ is simple.

Remember, a ring R being *simple* means:

$$\forall 0 \neq r \in R, \exists \alpha_i, \beta_i \in R \text{ with } \sum_{i=1}^n \alpha_i r \beta_i = 1_R.$$

Leavitt algebras

As a result, we have: Let S denote $L_K(1, n)$. Then

$$S^a \cong S^b \Leftrightarrow a \equiv b \pmod{n-1}.$$

In particular, $S \cong S^n$.

It turns out:

Theorem. (Leavitt, Duke J. Math, 1964)

For every field K and $n \geq 2$, $L_K(1, n)$ is simple.

Remember, a ring R being *simple* means:

$$\forall 0 \neq r \in R, \exists \alpha_i, \beta_i \in R \text{ with } \sum_{i=1}^n \alpha_i r \beta_i = 1_R.$$

Actually, $L_K(1, n)$ is REALLY simple:

$$\forall 0 \neq r \in L_K(1, n), \exists \alpha, \beta \in L_K(1, n) \text{ with } \alpha r \beta = 1_{L_K(1, n)}.$$

Building rings from combinatorial objects

Here's a familiar idea. Consider the set $T = \{x^0, x^1, x^2, \dots\}$. Define multiplication on T in the usual way: $x^i \cdot x^j = x^{i+j}$.

Consider formal symbols of the form

$$k_1 t_1 + k_2 t_2 + \cdots + k_n t_n$$

where $t_i \in T$, and $k_i \in \mathbb{R}$. Denote this set of symbols by KT . We can add and multiply two symbols of this form, as usual, e.g. $(kt)(k't') = kk'(t \cdot t')$.

Building rings from combinatorial objects

Here's a familiar idea. Consider the set $T = \{x^0, x^1, x^2, \dots\}$. Define multiplication on T in the usual way: $x^i \cdot x^j = x^{i+j}$.

Consider formal symbols of the form

$$k_1 t_1 + k_2 t_2 + \cdots + k_n t_n$$

where $t_i \in T$, and $k_i \in \mathbb{R}$. Denote this set of symbols by KT . We can add and multiply two symbols of this form, as usual, e.g. $(kt)(k't') = kk'(t \cdot t')$.

Here KT is just the ring $\mathbb{R}[x]$ of polynomials with coefficients in \mathbb{R} .

Building rings from combinatorial objects

Here's a familiar idea. Consider the set $T = \{x^0, x^1, x^2, \dots\}$. Define multiplication on T in the usual way: $x^i \cdot x^j = x^{i+j}$.

Consider formal symbols of the form

$$k_1 t_1 + k_2 t_2 + \cdots + k_n t_n$$

where $t_i \in T$, and $k_i \in \mathbb{R}$. Denote this set of symbols by KT . We can add and multiply two symbols of this form, as usual, e.g. $(kt)(k't') = kk'(t \cdot t')$.

Here KT is just the ring $\mathbb{R}[x]$ of polynomials with coefficients in \mathbb{R} .

Also, e.g. if we impose the relation $x^n = x^0$ on T , call the new semigroup \overline{T} , then $\overline{T} = \{x^0, x^1, x^2, \dots, x^{n-1}\}$, and

$$\mathbb{R}\overline{T} \cong \mathbb{R}[x]/\langle x^n - 1 \rangle$$

Building rings from combinatorial objects

This is a standard construction to produce rings:

Building rings from combinatorial objects

This is a standard construction to produce rings:

For instance:

matrix rings, group rings, multivariable polynomial rings, etc ...

General path algebras

Let E be a directed graph. (We will assume E is finite for this talk, but analysis can be done in general.) $E = (E^0, E^1, r, s)$

$$s(e) \bullet \xrightarrow{e} \bullet r(e)$$

The *path algebra of E with coefficients in K* is the K -algebra KS

$S =$ the set of all directed paths in E ,

multiplication of paths is juxtaposition. Denote by KE .

General path algebras

Let E be a directed graph. (We will assume E is finite for this talk, but analysis can be done in general.) $E = (E^0, E^1, r, s)$

$$s(e) \bullet \xrightarrow{e} \bullet r(e)$$

The *path algebra of E with coefficients in K* is the K -algebra KE

S = the set of all directed paths in E ,

multiplication of paths is juxtaposition. Denote by KE .

In particular, in KE ,

for each edge e , $s(e) \cdot e = e = e \cdot r(e)$

for each vertex v , $v \cdot v = v$

$$1_{KE} = \sum_{v \in E^0} v.$$


Building Leavitt path algebras

Start with E , build its *double graph* \widehat{E} .

Building Leavitt path algebras

Start with E , build its *double graph* \widehat{E} . Example:


$E =$




Building Leavitt path algebras

Start with E , build its *double graph* \widehat{E} . Example:

$E =$



$\widehat{E} =$



Building Leavitt path algebras

Construct the path algebra $K\widehat{E}$.

Building Leavitt path algebras

Construct the path algebra $K\widehat{E}$. Consider these relations in $K\widehat{E}$:

Building Leavitt path algebras

Construct the path algebra $K\widehat{E}$. Consider these relations in $K\widehat{E}$:

(CK1) $e^*e = r(e)$; and $f^*e = 0$ for $f \neq e$ (for all edges e, f in E).

(CK2) $v = \sum_{\{e \in E^1 \mid s(e)=v\}} ee^*$ for each vertex v in E .

Building Leavitt path algebras

Construct the path algebra $K\widehat{E}$. Consider these relations in $K\widehat{E}$:

(CK1) $e^*e = r(e)$; and $f^*e = 0$ for $f \neq e$ (for all edges e, f in E).

(CK2) $v = \sum_{\{e \in E^1 \mid s(e)=v\}} ee^*$ for each vertex v in E .

(just at those vertices v which are not *sinks*, and which emit only finitely many edges)

Building Leavitt path algebras

Construct the path algebra $K\widehat{E}$. Consider these relations in $K\widehat{E}$:

(CK1) $e^*e = r(e)$; and $f^*e = 0$ for $f \neq e$ (for all edges e, f in E).

(CK2) $v = \sum_{\{e \in E^1 \mid s(e)=v\}} ee^*$ for each vertex v in E .

(just at those vertices v which are not *sinks*, and which emit only finitely many edges)

Definition


The Leavitt path algebra of E with coefficients in K

$$L_K(E) = K\widehat{E} / \langle (CK1), (CK2) \rangle$$

Leavitt path algebras: Examples

Some sample computations in $L_{\mathbb{C}}(E)$ from the Example:

$\widehat{E} =$




$$ee^* + ff^* + gg^* = v \quad g^*g = w \quad g^*f = 0$$

$$h^*h = w \text{ (CK1)} \quad hh^* = u \text{ (CK2)}$$

Leavitt path algebras: Examples

Some sample computations in $L_{\mathbb{C}}(E)$ from the Example:

$\widehat{E} =$



$$ee^* + ff^* + gg^* = v \quad g^*g = w \quad g^*f = 0$$

$$h^*h = w \text{ (CK1)} \quad hh^* = u \text{ (CK2)}$$

$$ff^* = \dots \text{ (no simplification)} \quad \text{Note: } (ff^*)^2 = f(f^*f)f^* = ff^*$$

Leavitt path algebras: Examples

Standard algebras arising as Leavitt path algebras:

Leavitt path algebras: Examples

Standard algebras arising as Leavitt path algebras:

$$E = \bullet v_1 \xrightarrow{e_1} \bullet v_2 \xrightarrow{e_2} \bullet v_3 \dots \dots \dots \bullet v_{n-1} \xrightarrow{e_{n-1}} \bullet v_n$$

Then $L_K(E) \cong M_n(K)$.

Leavitt path algebras: Examples

Standard algebras arising as Leavitt path algebras:


$$E = \bullet v_1 \xrightarrow{e_1} \bullet v_2 \xrightarrow{e_2} \bullet v_3 \dots \bullet v_{n-1} \xrightarrow{e_{n-1}} \bullet v_n$$

Then $L_K(E) \cong M_n(K)$.

$$E = \bullet v \begin{array}{c} \curvearrowright \\ \curvearrowleft \end{array} x$$


Then $L_K(E) \cong K[x, x^{-1}]$.

Leavitt path algebras: Examples



Then $L_K(E) \cong L_K(1, n)$.

Leavitt path algebras: Examples




Then $L_K(E) \cong L_K(1, n)$.

$L_K(1, n)$ has generators and relations:

$$x_1, x_2, \dots, x_n, y_1, y_2, \dots, y_n \in L_K(1, n);$$

Leavitt path algebras: Examples




Then $L_K(E) \cong L_K(1, n)$.

$L_K(1, n)$ has generators and relations:

$$x_1, x_2, \dots, x_n, y_1, y_2, \dots, y_n \in L_K(1, n);$$

$$\sum_{i=1}^n y_i x_i = 1_{L_K(1, n)}, \quad \text{and} \quad x_i y_j = \delta_{i,j} 1_{L_K(1, n)},$$

Leavitt path algebras: Examples



Then $L_K(E) \cong L_K(1, n)$.

$L_K(1, n)$ has generators and relations:

$$x_1, x_2, \dots, x_n, y_1, y_2, \dots, y_n \in L_K(1, n);$$

$$\sum_{i=1}^n y_i x_i = 1_{L_K(1, n)}, \quad \text{and} \quad x_i y_j = \delta_{i,j} 1_{L_K(1, n)},$$

while $L_K(R_n)$ has these SAME generators and relations, where we identify y_i^* with x_i .

Historical note, part 1

1962: Leavitt gives construction of $L_K(1, n)$.

Historical note, part 1

1962: Leavitt gives construction of $L_K(1, n)$.

1979: Cuntz gives construction of the C^* -algebras \mathcal{O}_n .

Historical note, part 1

1962: Leavitt gives construction of $L_K(1, n)$.

1979: Cuntz gives construction of the C^* -algebras \mathcal{O}_n .

1980's: Cuntz, Krieger, and others generalize the \mathcal{O}_n construction to directed graphs, and produce the *graph C^* -algebras* $C^*(E)$.

Historical note, part 1

1962: Leavitt gives construction of $L_K(1, n)$.

1979: Cuntz gives construction of the C^* -algebras \mathcal{O}_n .

1980's: Cuntz, Krieger, and others generalize the \mathcal{O}_n construction to directed graphs, and produce the *graph C^* -algebras* $C^*(E)$.

June 2004: Various algebraists attend the CBMS lecture series

“Graph C^* -algebras: algebras we can see”,

held at University of Iowa, given by


Iain Raeburn (U. New South Wales / U. Newcastle / U. Wollongong ...);

algebraic analogs of graph C^* -algebras are defined and investigated starting Fall 2004.

- 1 Leavitt path algebras: Introduction and Motivation
- 2 Algebraic properties**
- 3 Projective modules
- 4 Connections and Applications


Some graph definitions

1. A *cycle*




Some graph definitions


1. A *cycle*



2. An *exit* for a cycle.



or



Some graph definitions

3a. *connects to* a vertex.


$$\bullet^u \longrightarrow \bullet^v \longrightarrow \bullet^w \quad (\text{also } \bullet^w)$$

Some graph definitions

3a. *connects to* a vertex.

$$\bullet^u \longrightarrow \bullet^v \longrightarrow \bullet^w \quad (\text{also } \bullet^w)$$

3b. *connects to* a cycle.



Simplicity of Leavitt path algebras

Here's a natural question, especially in light of Bill Leavitt's result that $L_K(1, n)$ is simple for all $n \geq 2$:

Simplicity of Leavitt path algebras

Here's a natural question, especially in light of Bill Leavitt's result that $L_K(1, n)$ is simple for all $n \geq 2$:

For which graphs E and fields K is $L_K(E)$ simple?

Simplicity of Leavitt path algebras


Here's a natural question, especially in light of Bill Leavitt's result that $L_K(1, n)$ is simple for all $n \geq 2$:

For which graphs E and fields K is $L_K(E)$ simple?

Note $L_K(E)$ is simple for

$$E = \bullet \longrightarrow \bullet \cdots \cdots \longrightarrow \bullet \quad \text{since } L_K(E) \cong M_n(K)$$

and for

and for $E = R_n =$  $\text{ since } L_K(E) \cong L_K(1, n)$

but not simple for

$$E = R_1 = \bullet \overset{v}{\curvearrowright} x \quad \text{since } L_K(E) \cong K[x, x^{-1}]$$

Simplicity of Leavitt path algebras

Theorem

(A -, Aranda Pino, 2005) $L_K(E)$ is simple if and only if:

- 1 Every vertex connects to every cycle and to every sink in E ,
and
- 2 Every cycle in E has an exit.

Simplicity of Leavitt path algebras

Theorem

(A -, Aranda Pino, 2005) $L_K(E)$ is simple if and only if:

- 1 Every vertex connects to every cycle and to every sink in E ,
and
- 2 Every cycle in E has an exit.

Note: No role played by K .

Other ring-theoretic properties of Leavitt path algebras

We know precisely the graphs E for which $L_K(E)$ has various other properties, e.g.:

- 1 one-sided chain conditions
- 2 prime
- 3 von Neumann regular
- 4 two-sided chain conditions
- 5 primitive

Many more.

- 1 Leavitt path algebras: Introduction and Motivation
- 2 Algebraic properties
- 3 Projective modules**
- 4 Connections and Applications

The monoid $\mathcal{V}(R)$

Recall: P is a *finitely generated projective* R -module in case
 $P \oplus Q \cong R^n$ for some Q , some $n \in \mathbb{N}$.

The monoid $\mathcal{V}(R)$

Recall: P is a *finitely generated projective* R -module in case
 $P \oplus Q \cong R^n$ for some Q , some $n \in \mathbb{N}$.

Key example: R itself, or any R^n .

The monoid $\mathcal{V}(R)$

Recall: P is a *finitely generated projective* R -module in case
 $P \oplus Q \cong R^n$ for some Q , some $n \in \mathbb{N}$.

Key example: R itself, or any R^n .

Additional examples: Rf where f is idempotent (i.e., $f^2 = f$),
since $Rf \oplus R(1 - f) = R^1$.

So, for example, in $R = M_2(\mathbb{R})$, $P = M_2(\mathbb{R})e_{1,1} = \begin{pmatrix} * & 0 \\ * & 0 \end{pmatrix}$
is a finitely projective R -module. Note $P \not\cong R^n$ for any n .

The monoid $\mathcal{V}(R)$

Recall: P is a *finitely generated projective* R -module in case
 $P \oplus Q \cong R^n$ for some Q , some $n \in \mathbb{N}$.

Key example: R itself, or any R^n .

Additional examples: Rf where f is idempotent (i.e., $f^2 = f$),
since $Rf \oplus R(1 - f) = R^1$.

So, for example, in $R = M_2(\mathbb{R})$, $P = M_2(\mathbb{R})e_{1,1} = \begin{pmatrix} * & 0 \\ * & 0 \end{pmatrix}$
is a finitely projective R -module. Note $P \not\cong R^n$ for any n .

So $L_K(E)$ contains projective modules of the form $L_K(E)ee^*$ for
each edge e of E .

The monoid $\mathcal{V}(R)$

$\mathcal{V}(R)$ denotes the isomorphism classes of finitely generated projective (left) R -modules. With operation \oplus , this becomes an abelian monoid. Note R itself plays a special role in $\mathcal{V}(R)$.

The monoid $\mathcal{V}(R)$

$\mathcal{V}(R)$ denotes the isomorphism classes of finitely generated projective (left) R -modules. With operation \oplus , this becomes an abelian monoid. Note R itself plays a special role in $\mathcal{V}(R)$.

Example. $R = K$, a field. Then $\mathcal{V}(R) \cong \mathbb{Z}^+$.

The monoid $\mathcal{V}(R)$

$\mathcal{V}(R)$ denotes the isomorphism classes of finitely generated projective (left) R -modules. With operation \oplus , this becomes an abelian monoid. Note R itself plays a special role in $\mathcal{V}(R)$.

Example. $R = K$, a field. Then $\mathcal{V}(R) \cong \mathbb{Z}^+$.

Example. $S = M_d(K)$, K a field. Then $\mathcal{V}(S) \cong \mathbb{Z}^+$.
(But note that the 'position' of S in $\mathcal{V}(S)$ is different than the position of R in $\mathcal{V}(R)$.)

The monoid $\mathcal{V}(R)$

$\mathcal{V}(R)$ denotes the isomorphism classes of finitely generated projective (left) R -modules. With operation \oplus , this becomes an abelian monoid. Note R itself plays a special role in $\mathcal{V}(R)$.

Example. $R = K$, a field. Then $\mathcal{V}(R) \cong \mathbb{Z}^+$.

Example. $S = M_d(K)$, K a field. Then $\mathcal{V}(S) \cong \mathbb{Z}^+$.
(But note that the 'position' of S in $\mathcal{V}(S)$ is different than the position of R in $\mathcal{V}(R)$.)

Remark: Given a ring R , it is in general not easy to compute $\mathcal{V}(R)$.

The monoid M_E

Here's a 'natural' monoid arising from any directed graph E .

The monoid M_E

Here's a 'natural' monoid arising from any directed graph E .

Associate to E the abelian monoid $(M_E, +)$:


$$M_E = \left\{ \sum_{v \in E^0} n_v a_v \right\}$$

with $n_v \in \mathbb{Z}^+$ for all $v \in E^0$.

Relations in M_E are given by: $a_v = \sum_{e \in s^{-1}(v)} a_r(e)$.

The monoid M_E


Example. Let F be the graph



So M_F consists of elements $\{n_1 a_1 + n_2 a_2 + n_3 a_3\}$ ($n_i \in \mathbb{Z}^+$),
subject to: $a_1 = a_2 + a_3$; $a_2 = a_1 + a_3$; $a_3 = a_1 + a_2$.

The monoid M_E

Example. Let F be the graph




So M_F consists of elements $\{n_1 a_1 + n_2 a_2 + n_3 a_3\}$ ($n_i \in \mathbb{Z}^+$),
subject to: $a_1 = a_2 + a_3$; $a_2 = a_1 + a_3$; $a_3 = a_1 + a_2$.

It's not hard to get:

The monoid M_E

Example. Let F be the graph




So M_F consists of elements $\{n_1 a_1 + n_2 a_2 + n_3 a_3\}$ ($n_i \in \mathbb{Z}^+$),
subject to: $a_1 = a_2 + a_3$; $a_2 = a_1 + a_3$; $a_3 = a_1 + a_2$.

It's not hard to get: $M_F = \{0, a_1, a_2, a_3, a_1 + a_2 + a_3\}$.

In particular, $M_F \setminus \{0\} \cong \mathbb{Z}_2 \times \mathbb{Z}_2$.

The monoid $\mathcal{V}(L_K(E))$

Example:




Then M_E is the set of symbols of the form

$$n_1 a_v \quad (n_1 \in \mathbb{Z}^+)$$

subject to the relation: $a_v = n a_v$

The monoid $\mathcal{V}(L_K(E))$

Example:



Then M_E is the set of symbols of the form

$$n_1 a_v \quad (n_1 \in \mathbb{Z}^+)$$

subject to the relation: $a_v = n a_v$

So here, $M_E = \{0, a_v, 2a_v, \dots, (n-1)a_v\}$.

In particular, $M_E \setminus \{0\} \cong \mathbb{Z}_{n-1}$.

The monoid $\mathcal{V}(L_K(E))$

The monoid $\mathcal{V}(L_K(E))$

Theorem

(P. Ara, M.A. Moreno, E. Pardo, 2007)

For any row-finite directed graph E ,

$$\mathcal{V}(L_K(E)) \cong M_E.$$

Moreover, $L_K(E)$ is universal with this property.

The monoid $\mathcal{V}(L_K(E))$

Theorem

(P. Ara, M.A. Moreno, E. Pardo, 2007)

For any row-finite directed graph E ,

$$\mathcal{V}(L_K(E)) \cong M_E.$$

Moreover, $L_K(E)$ is universal with this property.

One (very nontrivial) consequence: Let S denote $L_K(1, n)$. Then

$$\mathcal{V}(S) = \{0, S, S^2, \dots, S^{n-1}\}.$$

Historical Note, Part 2

So we can think of Leavitt path algebras in two ways:

- 1) the “quotient of a path algebra” approach, and
- 2) the “universal algebra which supports M_E as its \mathcal{V} -monoid” approach.

These were developed in parallel.

The two approaches together have complemented each other in the development of the subject.

Purely infinite simplicity

Here's a property (most likely unfamiliar to most of you ...)

Purely infinite simplicity

Here's a property (most likely unfamiliar to most of you ...)

We call a unital simple ring R *purely infinite simple* if R is not a division ring, and for every $r \neq 0$ in R there exists α, β in R for which

$$\alpha r \beta = 1_R.$$

Purely infinite simplicity

Leavitt showed that the Leavitt algebras $L_K(1, n)$ are in fact purely infinite simple.

Purely infinite simplicity

Leavitt showed that the Leavitt algebras $L_K(1, n)$ are in fact purely infinite simple.

Which Leavitt path algebras are purely infinite simple?

Purely infinite simplicity

Leavitt showed that the Leavitt algebras $L_K(1, n)$ are in fact purely infinite simple.

Which Leavitt path algebras are purely infinite simple?

Theorem:

$$L_K(E) \text{ is purely infinite simple} \Leftrightarrow$$
$$L_K(E) \text{ is simple,}$$

Purely infinite simplicity

Leavitt showed that the Leavitt algebras $L_K(1, n)$ are in fact purely infinite simple.

Which Leavitt path algebras are purely infinite simple?

Theorem:

$L_K(E)$ is purely infinite simple \Leftrightarrow
 $L_K(E)$ is simple, and E contains a cycle

Purely infinite simplicity

Leavitt showed that the Leavitt algebras $L_K(1, n)$ are in fact purely infinite simple.

Which Leavitt path algebras are purely infinite simple?

Theorem:

$L_K(E)$ is purely infinite simple \Leftrightarrow

$L_K(E)$ is simple, and E contains a cycle \Leftrightarrow

$M_E \setminus \{0\}$ is a group

Purely infinite simplicity

Leavitt showed that the Leavitt algebras $L_K(1, n)$ are in fact purely infinite simple.

Which Leavitt path algebras are purely infinite simple?

Theorem:

$L_K(E)$ is purely infinite simple \Leftrightarrow

$L_K(E)$ is simple, and E contains a cycle \Leftrightarrow

$M_E \setminus \{0\}$ is a group

Moreover, in this situation, we can easily calculate $\mathcal{V}(L_K(E))$ using the Smith normal form of the matrix $I - A_E$.

- 1 Leavitt path algebras: Introduction and Motivation
- 2 Algebraic properties
- 3 Projective modules
- 4 Connections and Applications**

Eight Year Update

In addition to “expected” types of results, over the past eight years Leavitt path algebras have played an interesting / important role in resolving various questions outside the subject per se.

- 1 Kaplansky’s question on prime non-primitive von Neumann regular algebras.
- 2 The realization question for von Neumann regular rings.
- 3 Constructing simple Lie algebras.
- 4 Connections to various C^* -algebras.
- 5 Constructing algebras with prescribed sets of prime / primitive ideals

Matrices over Leavitt algebras

One such connection:

Let $R = L_{\mathbb{C}}(1, n)$. So ${}_R R \cong {}_R R^n$.

So this gives in particular $R \cong M_n(R)$ as rings.

Which then (for free) gives some additional isomorphisms, e.g.

$$R \cong M_{n^i}(R)$$

for any $i \geq 1$.

Matrices over Leavitt algebras

One such connection:

Let $R = L_{\mathbb{C}}(1, n)$. So ${}_R R \cong {}_R R^n$.

So this gives in particular $R \cong M_n(R)$ as rings.

Which then (for free) gives some additional isomorphisms, e.g.

$$R \cong M_{n^i}(R)$$

for any $i \geq 1$.

Also, ${}_R R \cong {}_R R^n \cong {}_R R^{2n-1} \cong {}_R R^{3n-2} \cong \dots$, which also in turn yield ring isomorphisms

$$R \cong M_n(R) \cong M_{2n-1}(R) \cong M_{3n-2}(R) \cong \dots$$

Matrices over Leavitt algebras

Question: Are there other matrix sizes d for which $R \cong M_d(R)$?

Answer: In general, yes.

Matrices over Leavitt algebras

Question: Are there other matrix sizes d for which $R \cong M_d(R)$?

Answer: In general, yes.

For instance, if $R = L(1, 4)$, then it's not hard to show that $R \cong M_2(R)$ as rings (even though $R \not\cong_R R^2$ as modules).

Idea: 2 and 4 are nicely related, so these eight matrices inside $M_2(L(1, 4))$ “work”:

$$X_1 = \begin{pmatrix} x_1 & 0 \\ x_2 & 0 \end{pmatrix}, X_2 = \begin{pmatrix} x_3 & 0 \\ x_4 & 0 \end{pmatrix}, X_3 = \begin{pmatrix} 0 & x_1 \\ 0 & x_2 \end{pmatrix}, X_4 = \begin{pmatrix} 0 & x_3 \\ 0 & x_4 \end{pmatrix}$$

together with their duals

$$Y_1 = \begin{pmatrix} y_1 & y_2 \\ 0 & 0 \end{pmatrix}, Y_2 = \begin{pmatrix} y_3 & y_4 \\ 0 & 0 \end{pmatrix}, Y_3 = \begin{pmatrix} 0 & 0 \\ y_1 & y_2 \end{pmatrix}, Y_4 = \begin{pmatrix} 0 & 0 \\ y_3 & y_4 \end{pmatrix}$$

Matrices over Leavitt algebras

In general, using this same idea, we can show that:

if $d|n^t$ for some $t \in \mathbb{N}$, then $L(1, n) \cong M_d(L(1, n))$.

Matrices over Leavitt algebras

In general, using this same idea, we can show that:

if $d|n^t$ for some $t \in \mathbb{N}$, then $L(1, n) \cong M_d(L(1, n))$.

On the other hand ...

If $R = L(1, n)$, then the “type” of R is $n - 1$. (Think: “smallest difference”). Bill Leavitt showed the following in his 1962 paper:

The type of $M_d(L(1, n))$ is $\frac{n-1}{g.c.d.(d, n-1)}$.

In particular, if $g.c.d.(d, n - 1) > 1$, then $L(1, n) \not\cong M_d(L(1, n))$.

Matrices over Leavitt algebras

In general, using this same idea, we can show that:

if $d|n^t$ for some $t \in \mathbb{N}$, then $L(1, n) \cong M_d(L(1, n))$.

On the other hand ...

If $R = L(1, n)$, then the “type” of R is $n - 1$. (Think: “smallest difference”). Bill Leavitt showed the following in his 1962 paper:

The type of $M_d(L(1, n))$ is $\frac{n-1}{g.c.d.(d, n-1)}$.

In particular, if $g.c.d.(d, n - 1) > 1$, then $L(1, n) \not\cong M_d(L(1, n))$.

Conjecture: $L(1, n) \cong M_d(L(1, n)) \Leftrightarrow g.c.d.(d, n - 1) = 1$.

Matrices over Leavitt algebras

In general, using this same idea, we can show that:

if $d|n^t$ for some $t \in \mathbb{N}$, then $L(1, n) \cong M_d(L(1, n))$.

On the other hand ...

If $R = L(1, n)$, then the “type” of R is $n - 1$. (Think: “smallest difference”). Bill Leavitt showed the following in his 1962 paper:

The type of $M_d(L(1, n))$ is $\frac{n-1}{g.c.d.(d, n-1)}$.

In particular, if $g.c.d.(d, n - 1) > 1$, then $L(1, n) \not\cong M_d(L(1, n))$.

Conjecture: $L(1, n) \cong M_d(L(1, n)) \Leftrightarrow g.c.d.(d, n - 1) = 1$.

(Note: $d|n^t \Rightarrow g.c.d.(d, n - 1) = 1$.)

Matrices over Leavitt algebras

Smallest interesting pair: Is $L(1, 5) \cong M_3(L(1, 5))$?

We are led “naturally” to consider these five matrices (and their duals) in $M_3(L(1, 5))$:

$$\begin{pmatrix} x_1 & 0 & 0 \\ x_2 & 0 & 0 \\ x_3 & 0 & 0 \end{pmatrix} \quad \begin{pmatrix} x_4 & 0 & 0 \\ x_5 & 0 & 0 \\ 0 & 1 & 0 \end{pmatrix} \quad \begin{pmatrix} 0 & 0 & x_1^2 \\ 0 & 0 & x_2x_1 \\ 0 & 0 & x_3x_1 \end{pmatrix} \quad \begin{pmatrix} 0 & 0 & x_4x_1 \\ 0 & 0 & x_5x_1 \\ 0 & 0 & x_2 \end{pmatrix} \quad \begin{pmatrix} 0 & 0 & x_3 \\ 0 & 0 & x_4 \\ 0 & 0 & x_5 \end{pmatrix}$$

Everything went along swimmingly...

Matrices over Leavitt algebras

Smallest interesting pair: Is $L(1, 5) \cong M_3(L(1, 5))$?

We are led “naturally” to consider these five matrices (and their duals) in $M_3(L(1, 5))$:

$$\begin{pmatrix} x_1 & 0 & 0 \\ x_2 & 0 & 0 \\ x_3 & 0 & 0 \end{pmatrix} \quad \begin{pmatrix} x_4 & 0 & 0 \\ x_5 & 0 & 0 \\ 0 & 1 & 0 \end{pmatrix} \quad \begin{pmatrix} 0 & 0 & x_1^2 \\ 0 & 0 & x_2 x_1 \\ 0 & 0 & x_3 x_1 \end{pmatrix} \quad \begin{pmatrix} 0 & 0 & x_4 x_1 \\ 0 & 0 & x_5 x_1 \\ 0 & 0 & x_2 \end{pmatrix} \quad \begin{pmatrix} 0 & 0 & x_3 \\ 0 & 0 & x_4 \\ 0 & 0 & x_5 \end{pmatrix}$$

Everything went along swimmingly... But we couldn't see how to generate the matrix units $e_{1,3}$ and $e_{3,1}$ inside $M_3(L(1, 5))$ using these ten matrices.

Matrices over Leavitt algebras

Breakthrough (came from an analysis of isomorphisms between more general Leavitt path algebras) ... we were using the wrong ten matrices.

Matrices over Leavitt algebras

Breakthrough (came from an analysis of isomorphisms between more general Leavitt path algebras) ... we were using the wrong ten matrices. Original set:

$$\begin{pmatrix} x_1 & 0 & 0 \\ x_2 & 0 & 0 \\ x_3 & 0 & 0 \end{pmatrix} \quad \begin{pmatrix} x_4 & 0 & 0 \\ x_5 & 0 & 0 \\ 0 & 1 & 0 \end{pmatrix} \quad \begin{pmatrix} 0 & 0 & x_1^2 \\ 0 & 0 & x_2x_1 \\ 0 & 0 & x_3x_1 \end{pmatrix} \quad \begin{pmatrix} 0 & 0 & x_4x_1 \\ 0 & 0 & x_5x_1 \\ 0 & 0 & x_2 \end{pmatrix} \quad \begin{pmatrix} 0 & 0 & x_3 \\ 0 & 0 & x_4 \\ 0 & 0 & x_5 \end{pmatrix}$$

Matrices over Leavitt algebras

Breakthrough (came from an analysis of isomorphisms between more general Leavitt path algebras) ... we were using the wrong ten matrices. Original set:

$$\begin{pmatrix} x_1 & 0 & 0 \\ x_2 & 0 & 0 \\ x_3 & 0 & 0 \end{pmatrix} \quad \begin{pmatrix} x_4 & 0 & 0 \\ x_5 & 0 & 0 \\ 0 & 1 & 0 \end{pmatrix} \quad \begin{pmatrix} 0 & 0 & x_1^2 \\ 0 & 0 & x_2x_1 \\ 0 & 0 & x_3x_1 \end{pmatrix} \quad \begin{pmatrix} 0 & 0 & x_4x_1 \\ 0 & 0 & x_5x_1 \\ 0 & 0 & x_2 \end{pmatrix} \quad \begin{pmatrix} 0 & 0 & x_3 \\ 0 & 0 & x_4 \\ 0 & 0 & x_5 \end{pmatrix}$$

Instead, this set (together with duals) works:

$$\begin{pmatrix} x_1 & 0 & 0 \\ x_2 & 0 & 0 \\ x_3 & 0 & 0 \end{pmatrix} \quad \begin{pmatrix} x_4 & 0 & 0 \\ x_5 & 0 & 0 \\ 0 & 1 & 0 \end{pmatrix} \quad \begin{pmatrix} 0 & 0 & x_1^2 \\ 0 & 0 & x_2x_1 \\ 0 & 0 & x_3x_1 \end{pmatrix} \quad \begin{pmatrix} 0 & 0 & x_4x_1 \\ 0 & 0 & x_5x_1 \\ 0 & 0 & x_2 \end{pmatrix} \quad \begin{pmatrix} 0 & 0 & x_4 \\ 0 & 0 & x_3 \\ 0 & 0 & x_5 \end{pmatrix}$$

Matrices over Leavitt algebras

Theorem

(A-, Ánh, Pardo; Crelle's J. 2008) For any field K ,

$$L_K(1, n) \cong M_d(L_K(1, n)) \Leftrightarrow \text{g.c.d.}(d, n-1) = 1.$$

Indeed, more generally,

$$M_d(L_K(1, n)) \cong M_{d'}(L_K(1, n')) \Leftrightarrow \\ n = n' \text{ and } \text{g.c.d.}(d, n-1) = \text{g.c.d.}(d', n-1).$$

Moreover, we can write down the isomorphisms explicitly.

Matrices over Leavitt algebras

Theorem

(A-, Ánh, Pardo; *Crelle's J.* 2008) For any field K ,

$$L_K(1, n) \cong M_d(L_K(1, n)) \Leftrightarrow \text{g.c.d.}(d, n-1) = 1.$$

Indeed, more generally,

$$M_d(L_K(1, n)) \cong M_{d'}(L_K(1, n')) \Leftrightarrow \\ n = n' \text{ and } \text{g.c.d.}(d, n-1) = \text{g.c.d.}(d', n-1).$$

Moreover, we can write down the isomorphisms explicitly.

Along the way, some elementary (but apparently new) number theory ideas come into play.

Given n, d with $\text{g.c.d.}(d, n - 1) = 1$, there is a “natural” partition of $\{1, 2, \dots, n\}$ into two disjoint subsets.

Given n, d with $\text{g.c.d.}(d, n - 1) = 1$, there is a “natural” partition of $\{1, 2, \dots, n\}$ into two disjoint subsets.

Here’s what made this second set of matrices work. Using this partition in the particular case $n = 5, d = 3$, then the partition of $\{1, 2, 3, 4, 5\}$ turns out to be the two sets

$$\{1, 4\} \quad \text{and} \quad \{2, 3, 5\}.$$

The matrices that “worked” are ones where we fill in the last columns with terms of the form $x_i x_1^j$ in such a way that i is in the same subset as the row number of that entry.

Given n, d with $\text{g.c.d.}(d, n - 1) = 1$, there is a “natural” partition of $\{1, 2, \dots, n\}$ into two disjoint subsets.

Here's what made this second set of matrices work. Using this partition in the particular case $n = 5, d = 3$, then the partition of $\{1, 2, 3, 4, 5\}$ turns out to be the two sets

$$\{1, 4\} \quad \text{and} \quad \{2, 3, 5\}.$$

The matrices that “worked” are ones where we fill in the last columns with terms of the form $x_i x_1^j$ in such a way that i is in the same subset as the row number of that entry.

The number theory underlying this partition in the general case where $\text{g.c.d.}(d, n - 1) = 1$ is elementary. But we are hoping to find some other 'context' in which this partition process arises.

Matrices over Leavitt algebras

Computations when $n = 5, d = 3$.

$\gcd(3, 5 - 1) = 1$. Now $5 = 1 \cdot 3 + 2$, so that $r = 2, r - 1 = 1$, and define $s = d - (r - 1) = 3 - 1 = 2$.

Consider the sequence starting at 1, and increasing by s each step, and interpret mod d ($1 \leq i \leq d$). This will necessarily give all integers between 1 and d .

Matrices over Leavitt algebras

Computations when $n = 5, d = 3$.

$\gcd(3, 5 - 1) = 1$. Now $5 = 1 \cdot 3 + 2$, so that $r = 2, r - 1 = 1$, and define $s = d - (r - 1) = 3 - 1 = 2$.

Consider the sequence starting at 1, and increasing by s each step, and interpret mod d ($1 \leq i \leq d$). This will necessarily give all integers between 1 and d .

So here we get the sequence $1, 3, 2$.

Matrices over Leavitt algebras

Computations when $n = 5, d = 3$.

$\gcd(3, 5 - 1) = 1$. Now $5 = 1 \cdot 3 + 2$, so that $r = 2, r - 1 = 1$, and define $s = d - (r - 1) = 3 - 1 = 2$.

Consider the sequence starting at 1, and increasing by s each step, and interpret mod d ($1 \leq i \leq d$). This will necessarily give all integers between 1 and d .

So here we get the sequence $1, 3, 2$.

Now break this set into two pieces: those integers up to and including $r - 1$, and those after. Since $r - 1 = 1$, here we get

$$\{1, 2, 3\} = \{1\} \cup \{2, 3\}.$$

Matrices over Leavitt algebras

Computations when $n = 5, d = 3$.

$\gcd(3, 5 - 1) = 1$. Now $5 = 1 \cdot 3 + 2$, so that $r = 2, r - 1 = 1$, and define $s = d - (r - 1) = 3 - 1 = 2$.

Consider the sequence starting at 1, and increasing by s each step, and interpret mod d ($1 \leq i \leq d$). This will necessarily give all integers between 1 and d .

So here we get the sequence $1, 3, 2$.

Now break this set into two pieces: those integers up to and including $r - 1$, and those after. Since $r - 1 = 1$, here we get

$$\{1, 2, 3\} = \{1\} \cup \{2, 3\}.$$

Now extend these two sets mod 3 to all integers up to 5.

$$\{1, 4\} \cup \{2, 3, 5\}$$



Matrices over Leavitt algebras

Does this look familiar?

Matrices over Leavitt algebras

Corollary. (Matrices over the Cuntz C^* -algebras)

$$\mathcal{O}_n \cong M_d(\mathcal{O}_n) \Leftrightarrow g.c.d.(d, n-1) = 1.$$

(And the isomorphisms are explicitly described.)

Matrices over Leavitt algebras

An important recent application:

For each pair of positive integers n, r , there exists an infinite, finitely presented simple group $G_{n,r}^+$. These were introduced in:

G. Higman, “Finitely presented infinite simple groups”, Notes on Pure Mathematics, 8, Department of Pure Mathematics, I.A.S. Australian National University, Canberra, 1974.

Matrices over Leavitt algebras

An important recent application:

For each pair of positive integers n, r , there exists an infinite, finitely presented simple group $G_{n,r}^+$. These were introduced in:

G. Higman, “Finitely presented infinite simple groups”, Notes on Pure Mathematics, 8, Department of Pure Mathematics, I.A.S. Australian National University, Canberra, 1974.

Theorem. (E. Pardo, 2011)

$$G_{n,r}^+ \cong G_{m,s}^+ \iff m = n \text{ and } \text{g.c.d.}(r, n - 1) = \text{g.c.d.}(s, n - 1).$$

Matrices over Leavitt algebras

An important recent application:

For each pair of positive integers n, r , there exists an infinite, finitely presented simple group $G_{n,r}^+$. These were introduced in:

G. Higman, “Finitely presented infinite simple groups”, Notes on Pure Mathematics, 8, Department of Pure Mathematics, I.A.S. Australian National University, Canberra, 1974.

Theorem. (E. Pardo, 2011)

$$G_{n,r}^+ \cong G_{m,s}^+ \iff m = n \text{ and } \text{g.c.d.}(r, n - 1) = \text{g.c.d.}(s, n - 1).$$

Proof. Show that $G_{n,r}^+$ can be realized as an appropriate subgroup of the invertible elements of $M_r(L_{\mathbb{C}}(1, n))$, and then use the explicit isomorphisms provided in the A - , Ánh, Pardo result.

What else is out there?

What else is out there?

$$(1) \quad L_K(E) \cong L_K(F) \Leftrightarrow \text{???}$$

Ideas from symbolic dynamics come into play here. Using some results on flow equivalence, we have been able to show:

Theorem. (A -, Louly, Pardo, Smith, 2011) If $L_K(E)$ and $L_K(F)$ are purely infinite simple Leavitt path algebras such that

$$(K_0(L_K(E)), [1_{L_K(E)}]) \cong (K_0(L_K(F)), [1_{L_K(F)}]),$$

What else is out there?

$$(1) \quad L_K(E) \cong L_K(F) \Leftrightarrow ???$$

Ideas from symbolic dynamics come into play here. Using some results on flow equivalence, we have been able to show:

Theorem. (A -, Louly, Pardo, Smith, 2011) If $L_K(E)$ and $L_K(F)$ are purely infinite simple Leavitt path algebras such that

$$(K_0(L_K(E)), [1_{L_K(E)}]) \cong (K_0(L_K(F)), [1_{L_K(F)}]),$$

$$\text{and } \det(I - A_E^t) = \det(I - A_F^t),$$

then $L_K(E) \cong L_K(F)$.

What else is out there?

$$(1) \quad L_K(E) \cong L_K(F) \Leftrightarrow \quad ? \quad ? \quad ?$$

Ideas from symbolic dynamics come into play here. Using some results on flow equivalence, we have been able to show:

Theorem. (A -, Louly, Pardo, Smith, 2011) If $L_K(E)$ and $L_K(F)$ are purely infinite simple Leavitt path algebras such that

$$(K_0(L_K(E)), [1_{L_K(E)}]) \cong (K_0(L_K(F)), [1_{L_K(F)}]),$$

$$\text{and } \det(I - A_E^t) = \det(I - A_F^t),$$

then $L_K(E) \cong L_K(F)$. Can we drop the determinant hypothesis?

What else is out there?

In particular, if



is $L_{\mathbb{C}}(E_4) \cong L_{\mathbb{C}}(1, 2)$?

What else is out there?

In particular, if



is $L_{\mathbb{C}}(E_4) \cong L_{\mathbb{C}}(1, 2)$?

The answer will be interesting, however it plays out.

What else is out there?

(2) For any graph E there is an intimate relationship between $L_{\mathbb{C}}(E)$ and $C^*(E)$. There are many theorems of the form:

$L_{\mathbb{C}}(E)$ has algebraic property $\mathcal{P} \Leftrightarrow C^*(E)$ has analytic property \mathcal{P}

What else is out there?

(2) For any graph E there is an intimate relationship between $L_{\mathbb{C}}(E)$ and $C^*(E)$. There are many theorems of the form:

$L_{\mathbb{C}}(E)$ has algebraic property $\mathcal{P} \Leftrightarrow C^*(E)$ has analytic property \mathcal{P}

but the proofs are not direct! They all are based on showing that the two properties are both equivalent to

E has graph property \mathcal{Q} .

Why this happens is still a mystery.

What else is out there?

Note: Questions (1) and (2) will take up much of the discussion at:

“The AMSI Workshop on Graph C^* -algebras,
Leavitt path algebras and symbolic dynamics”

February 11 - 14 University of Western Sydney

www.amsi.org.au → Research → Workshop Programs

What else is out there?

(3) Mad Veterinarian puzzles!

What else is out there?

(3) Mad Veterinarian puzzles!

(A-, Jessica Sklar), “The graph menagerie: Abstract algebra meets the Mad Veterinarian”, *Mathematics Magazine* **83**(3), 2010, 168-179.

Questions?

Thanks to:

Australian Mathematical Sciences Institute

Australian Mathematical Society

University of Western Sydney

Simons Foundation

Some elementary number theory

The partition of $\{1, 2, \dots, n\}$ induced by d when $\text{g.c.d.}(d, n - 1) = 1$

Suppose $\text{g.c.d.}(d, n - 1) = 1$. Write

$$n = dt + r \text{ with } 1 \leq r \leq d.$$

Let s denote $d - (r - 1)$.

It is easy to show that $\text{g.c.d.}(d, n - 1) = 1$ implies $\text{g.c.d.}(d, s) = 1$. We consider the sequence $\{h_i\}_{i=1}^d$ of integers, whose i^{th} entry is given by

$$h_i = 1 + (i - 1)s \pmod{d}.$$

Some elementary number theory

The integers h_i are understood to be taken from the set $\{1, 2, \dots, d\}$.

Because $\text{g.c.d.}(d, s) = 1$, basic number theory yields that the set of entries $\{h_1, h_2, \dots, h_d\}$ equals the set $\{1, 2, \dots, d\}$ (in some order). Our interest lies in a decomposition of $\{1, 2, \dots, d\}$ effected by the sequence h_1, h_2, \dots, h_d , as follows.

Some elementary number theory

We let d_1 denote the integer for which

$$h_{d_1} = r - 1$$

in the previously defined sequence. We denote by \hat{S}_1 the following subset of $\{1, 2, \dots, d\}$:

$$\hat{S}_1 = \{h_i \mid 1 \leq i \leq d_1\}.$$

We denote by \hat{S}_2 the complement of \hat{S}_1 in $\{1, 2, \dots, d\}$. We now construct a partition $S_1 \cup S_2$ of $\{1, 2, \dots, n\}$ by defining, for each $j \in \{1, 2, \dots, n\}$ and for $i \in \{1, 2\}$,

$j \in S_i$ precisely when $j \equiv j' \pmod{d}$ for $j' \in \{1, 2, \dots, d\}$, and $j' \in \hat{S}_i$.

(In other words, we extend the partition $\hat{S}_1 \cup \hat{S}_2$ of $\{1, 2, \dots, d\}$ to a partition $S_1 \cup S_2$ of $\{1, 2, \dots, n\}$ by extending mod d .)

Some elementary number theory

Example. Suppose $n = 35$, $d = 13$. Then $\gcd(13, 35 - 1) = 1$, so we are in the desired situation. Now $35 = 2 \cdot 13 + 9$, so that $r = 9$, $r - 1 = 8$, and $s = d - (r - 1) = 13 - 8 = 5$. Then we consider the sequence starting at 1, and increasing by s each step, and interpret mod d . (This will give all integers between 1 and d .)

Some elementary number theory

Example. Suppose $n = 35$, $d = 13$. Then $\gcd(13, 35 - 1) = 1$, so we are in the desired situation. Now $35 = 2 \cdot 13 + 9$, so that $r = 9$, $r - 1 = 8$, and $s = d - (r - 1) = 13 - 8 = 5$. Then we consider the sequence starting at 1, and increasing by s each step, and interpret mod d . (This will give all integers between 1 and d .) So here we get the sequence 1, 6, 11, 3, 8, 13, 5, 10, 2, 7, 12, 4, 9.

Some elementary number theory

Example. Suppose $n = 35$, $d = 13$. Then $\gcd(13, 35 - 1) = 1$, so we are in the desired situation. Now $35 = 2 \cdot 13 + 9$, so that $r = 9$, $r - 1 = 8$, and $s = d - (r - 1) = 13 - 8 = 5$. Then we consider the sequence starting at 1, and increasing by s each step, and interpret mod d . (This will give all integers between 1 and d .)

So here we get the sequence 1, 6, 11, 3, 8, 13, 5, 10, 2, 7, 12, 4, 9.

Now break this set into two pieces: those integers up to and including $r - 1$, and those after. Since $r - 1 = 8$, here we get

$$\{1, 2, \dots, 13\} = \{1, 3, 6, 8, 11\} \cup \{2, 4, 5, 7, 9, 10, 12, 13\}.$$

Some elementary number theory

Example. Suppose $n = 35$, $d = 13$. Then $\gcd(13, 35 - 1) = 1$, so we are in the desired situation. Now $35 = 2 \cdot 13 + 9$, so that $r = 9$, $r - 1 = 8$, and $s = d - (r - 1) = 13 - 8 = 5$. Then we consider the sequence starting at 1, and increasing by s each step, and interpret mod d . (This will give all integers between 1 and d .)

So here we get the sequence 1, 6, 11, 3, 8, 13, 5, 10, 2, 7, 12, 4, 9.

Now break this set into two pieces: those integers up to and including $r - 1$, and those after. Since $r - 1 = 8$, here we get

$$\{1, 2, \dots, 13\} = \{1, 3, 6, 8, 11\} \cup \{2, 4, 5, 7, 9, 10, 12, 13\}.$$

Now extend these two sets mod 13 to all integers up to 35.

$$\{1, 3, 6, 8, 11, 14, 16, 19, 21, 24, 27, 29, 32, 34\} \cup \\ \{2, 4, 5, 7, 9, 10, 12, 13, 15, 17, 18, 20, 22, 23, 25, 26, 28, 30, 31, 33, 35\}$$

