

Primitive graph algebras

Gene Abrams



BIRS Workshop: “Graph algebras: Bridges between graph C^* -algebras and Leavitt path algebras”

April 2013

Overview

1 Primitive Leavitt path algebras

2 Primitive graph C^* -algebras

1 Primitive Leavitt path algebras

2 Primitive graph C^* -algebras

Throughout R is associative, but not necessarily with identity.

Assume R at least has “local units”:

Prime rings

Definition: I, J two-sided ideals of R . The product IJ is the two-sided ideal

$$IJ = \left\{ \sum_{\ell=1}^n i_{\ell} j_{\ell} \mid i_{\ell} \in I, j_{\ell} \in J, n \in \mathbb{N} \right\}.$$

R is *prime* if the product of any two nonzero two-sided ideals of R is nonzero.

Prime rings

Definition: I, J two-sided ideals of R . The product IJ is the two-sided ideal

$$IJ = \left\{ \sum_{\ell=1}^n i_{\ell} j_{\ell} \mid i_{\ell} \in I, j_{\ell} \in J, n \in \mathbb{N} \right\}.$$

R is *prime* if the product of any two nonzero two-sided ideals of R is nonzero.

Examples:

- 1 Commutative domains, e.g. fields, \mathbb{Z} , $K[x]$, $K[x, x^{-1}]$, ...
- 2 Simple rings
- 3 $\text{End}_K(V)$ where $\dim_K(V)$ is infinite. $(\cong \text{RFM}(K))$

Prime rings

Note: Definition makes sense for nonunital rings.

Prime rings

Note: Definition makes sense for nonunital rings.

Lemma: R prime. Then R embeds as an ideal in a unital prime ring R_1 . (Dorroh extension of R .)

If R is a K -algebra then we can construct R_1 a K -algebra for which $\dim_K(R_1/R) = 1$.

Primitive rings

Definition: R is *left primitive* if R admits a faithful simple (= “irreducible”) left R -module.

Rephrased: if there exists ${}_R M$ simple for which $\text{Ann}_R(M) = \{0\}$.

Primitive rings

Definition: R is *left primitive* if R admits a faithful simple (= “irreducible”) left R -module.

Rephrased: if there exists ${}_R M$ simple for which $\text{Ann}_R(M) = \{0\}$.

Examples:

- Simple rings (note: need local units to build irreducibles)

NON-Examples:

- \mathbb{Z} , $K[x]$, $K[x, x^{-1}]$

Primitive rings

Primitive rings are “natural” generalizations of matrix rings.

Jacobson’s Density Theorem: R is primitive if and only if R is isomorphic to a dense subring of $\text{End}_D(V)$, for some division ring D , and some D -vector space V .

Primitive rings

Primitive rings are “natural” generalizations of matrix rings.

Jacobson’s Density Theorem: R is primitive if and only if R is isomorphic to a dense subring of $\text{End}_D(V)$, for some division ring D , and some D -vector space V .

Here $D = \text{End}_R(M)$ where M is the supposed simple faithful R -module.

Primitive rings

Primitive rings are “natural” generalizations of matrix rings.

Jacobson’s Density Theorem: R is primitive if and only if R is isomorphic to a dense subring of $\text{End}_D(V)$, for some division ring D , and some D -vector space V .

Here $D = \text{End}_R(M)$ where M is the supposed simple faithful R -module.

So this gives many more examples of primitive rings, e.g. $\text{FM}(K)$, $\text{RCFM}(K)$, etc ...

Primitive rings

Primitive rings are “natural” generalizations of matrix rings.

Jacobson’s Density Theorem: R is primitive if and only if R is isomorphic to a dense subring of $\text{End}_D(V)$, for some division ring D , and some D -vector space V .

Here $D = \text{End}_R(M)$ where M is the supposed simple faithful R -module.

So this gives many more examples of primitive rings, e.g. $\text{FM}(K)$, $\text{RCFM}(K)$, etc ...

Definition of “primitive” makes sense for non-unital rings.

Prime and primitive rings

Lemma: Every primitive ring is prime.

Prime and primitive rings

Lemma: Every primitive ring is prime.

Proof. Let M denote a simple faithful left R -module. Suppose $I \cdot J = \{0\}$. We want to show either $I = \{0\}$ or $J = \{0\}$.

Prime and primitive rings

Lemma: Every primitive ring is prime.

Proof. Let M denote a simple faithful left R -module. Suppose $I \cdot J = \{0\}$. We want to show either $I = \{0\}$ or $J = \{0\}$.

So $(I \cdot J)M = 0$. If $JM = \{0\}$ then $J = \{0\}$ as M is faithful. So suppose $JM \neq 0$. Then $JM = M$ (as M is simple), so $(I \cdot J)M = 0$ gives $IM = 0$, so $I = \{0\}$ as M is faithful. \square

Prime and primitive rings

Lemma: Every primitive ring is prime.

Proof. Let M denote a simple faithful left R -module. Suppose $I \cdot J = \{0\}$. We want to show either $I = \{0\}$ or $J = \{0\}$.

So $(I \cdot J)M = 0$. If $JM = \{0\}$ then $J = \{0\}$ as M is faithful. So suppose $JM \neq 0$. Then $JM = M$ (as M is simple), so $(I \cdot J)M = 0$ gives $IM = 0$, so $I = \{0\}$ as M is faithful. \square

If R is prime, then in previous embedding,

$$R \text{ is primitive} \Leftrightarrow R_1 \text{ is primitive.}$$

Prime and primitive rings

Converse of Lemma is not true (e.g. \mathbb{Z} , $K[x]$, $K[x, x^{-1}]$).

In fact, the only commutative primitive unital rings are fields.

Prime and primitive rings

Converse of Lemma is not true (e.g. \mathbb{Z} , $K[x]$, $K[x, x^{-1}]$).

In fact, the only commutative primitive unital rings are fields.

Remark for later:

From a ring-theoretic point of view, the question of finding prime, non-primitive rings is uninteresting (since there are so many of them!)

Prime and primitive rings

Remark: For ${}_R M$ simple, write $M \cong R/N$ for N a maximal left ideal of R . How can $\text{Ann}_R(M) = \{0\}$?

Prime and primitive rings

Remark: For ${}_R M$ simple, write $M \cong R/N$ for N a maximal left ideal of R . How can $\text{Ann}_R(M) = \{0\}$?

Note $n \cdot (r + N) = nr + N$ need not be $\bar{0}$ in R/N since nr is not necessarily in N .

Example: K any field, V an infinite dimensional K -vector space. $R = \text{End}_K(V) \cong \text{RFM}(K)$ is primitive, not simple.

Here $M = Re_{11}$ is simple. Easy to show $\text{Ann}_R(M) = \{0\}$, but R contains a nontrivial ideal (the finite rank transformations).

Prime and primitive rings

Remark: For ${}_R M$ simple, write $M \cong R/N$ for N a maximal left ideal of R . How can $\text{Ann}_R(M) = \{0\}$?

Note $n \cdot (r + N) = nr + N$ need not be $\bar{0}$ in R/N since nr is not necessarily in N .

Example: K any field, V an infinite dimensional K -vector space. $R = \text{End}_K(V) \cong \text{RFM}(K)$ is primitive, not simple.

Here $M = Re_{11}$ is simple. Easy to show $\text{Ann}_R(M) = \{0\}$, but R contains a nontrivial ideal (the finite rank transformations).

But we always have $\text{Ann}_R(R/N) \subseteq N$, since if $r(1 + N) = 0 + N$ then $r \in N$.

Leavitt path algebras

Let K be a field, and let $E = (E^0, E^1, s, r)$ be **any** directed graph.

The Leavitt path K -algebra $L_K(E)$ of E with coefficients in K

is the K -algebra generated by a set $\{v \mid v \in E^0\}$, together with a set of variables $\{e, e^* \mid e \in E^1\}$, which satisfy the following relations:

$$(V) \quad vw = \delta_{v,w}v \text{ for all } v, w \in E^0,$$

$$(E1) \quad s(e)e = er(e) = e \text{ for all } e \in E^1,$$

$$(E2) \quad r(e)e^* = e^*s(e) = e^* \text{ for all } e \in E^1, \text{ and}$$

$$(CK1) \quad e^*e' = \delta_{e,e'}r(e) \text{ for all } e, e' \in E^1.$$

$$(CK2) \quad v = \sum_{\{e \in E^1 \mid s(e)=v\}} ee^* \text{ for every regular vertex } v \in E^0.$$

Prime Leavitt path algebras

Notation: $u \geq v$ means either $u = v$ or there exists a path p for which $s(p) = u, r(p) = v$. u “connects to” v .

Prime Leavitt path algebras

Notation: $u \geq v$ means either $u = v$ or there exists a path p for which $s(p) = u, r(p) = v$. u “connects to” v .

Theorem. (Aranda Pino, Pardo, Siles Molina 2009) E arbitrary. Then $L_K(E)$ is prime \Leftrightarrow for each pair $v, w \in E^0$ there exists $u \in E^0$ with $v \geq u$ and $w \geq u$.

Prime Leavitt path algebras

Notation: $u \geq v$ means either $u = v$ or there exists a path p for which $s(p) = u, r(p) = v$. u “connects to” v .

Theorem. (Aranda Pino, Pardo, Siles Molina 2009) E arbitrary. Then $L_K(E)$ is prime \Leftrightarrow for each pair $v, w \in E^0$ there exists $u \in E^0$ with $v \geq u$ and $w \geq u$. “Downward Directed” (MT3)

Prime Leavitt path algebras

Notation: $u \geq v$ means either $u = v$ or there exists a path p for which $s(p) = u, r(p) = v$. u “connects to” v .

Theorem. (Aranda Pino, Pardo, Siles Molina 2009) E arbitrary. Then $L_K(E)$ is prime \Leftrightarrow for each pair $v, w \in E^0$ there exists $u \in E^0$ with $v \geq u$ and $w \geq u$. “Downward Directed” (MT3)

Idea of Proof. (\Rightarrow) Let R denote $L_K(E)$. Let $v, w \in E^0$. But $RvR \neq \{0\}$ and $RwR \neq \{0\} \Rightarrow RvRwR \neq \{0\} \Rightarrow vRw \neq \{0\} \Rightarrow v\alpha\beta^*w \neq 0$ for some paths α, β in E . Then $u = r(\alpha)$ works.

Prime Leavitt path algebras

Notation: $u \geq v$ means either $u = v$ or there exists a path p for which $s(p) = u, r(p) = v$. u “connects to” v .

Theorem. (Aranda Pino, Pardo, Siles Molina 2009) E arbitrary. Then $L_K(E)$ is prime \Leftrightarrow for each pair $v, w \in E^0$ there exists $u \in E^0$ with $v \geq u$ and $w \geq u$. “Downward Directed” (MT3)

Idea of Proof. (\Rightarrow) Let R denote $L_K(E)$. Let $v, w \in E^0$. But $RvR \neq \{0\}$ and $RwR \neq \{0\} \Rightarrow RvRwR \neq \{0\} \Rightarrow vRw \neq \{0\} \Rightarrow v\alpha\beta^*w \neq 0$ for some paths α, β in E . Then $u = r(\alpha)$ works.

(\Leftarrow) $L_K(E)$ is graded by \mathbb{Z} , so need only check primeness on nonzero graded ideals I, J . But each nonzero graded ideal contains a vertex. Let $v \in I \cap E^0$ and $w \in J \cap E^0$. By downward directedness there is $u \in E^0$ with $v \geq u$ and $w \geq u$. But then $u = p^*vp \in I$ and $u = q^*wq \in J$, so that $0 \neq u = u^2 \in IJ$.

The Countable Separation Property

‘ **Definition.** Let E be any directed graph. E has the *Countable Separation Property* (CSP) if there exists a countable set of vertices S in E for which every vertex of E connects to an element of S .

E has the “Countable Separation Property” with respect to S .

The Countable Separation Property

‘ **Definition.** Let E be any directed graph. E has the *Countable Separation Property* (CSP) if there exists a countable set of vertices S in E for which every vertex of E connects to an element of S .

E has the “Countable Separation Property” with respect to S .

Same idea for any subset X of E^0 : X has CSP (with respect to S_X) in case S_X is countable, and every element of X connects to an element of S_X .

The Countable Separation Property

Definition. Let E be any directed graph. E has the *Countable Separation Property* (CSP) if there exists a countable set of vertices S in E for which every vertex of E connects to an element of S .

E has the “Countable Separation Property” with respect to S .

Same idea for any subset X of E^0 : X has CSP (with respect to S_X) in case S_X is countable, and every element of X connects to an element of S_X .

Note for later: If $X = \emptyset$, then X vacuously has CSP (with respect to any countable subset of vertices).

So if X does not have CSP, then $X \neq \emptyset$.

The Countable Separation Property

Observe: If E^0 is countable, then E has CSP.

The Countable Separation Property

Observe: If E^0 is countable, then E has CSP.

2) **Example:** X uncountable, S the set of finite subsets of X .
Define the graph E :

- 1 vertices indexed by S , and
- 2 edges induced by proper subset relationship.

Then E does not have CSP.

Primitive Leavitt path algebras

Can we describe the (left) primitive Leavitt path algebras?

Note: Since $L_K(E) \cong L_K(E)^{op}$, left primitivity and right primitivity coincide. So we can just say “primitive” Leavitt path algebra.

Primitive Leavitt path algebras

Can we describe the (left) primitive Leavitt path algebras?

Note: Since $L_K(E) \cong L_K(E)^{op}$, left primitivity and right primitivity coincide. So we can just say “primitive” Leavitt path algebra.

Theorem. (A-, Jason Bell, K.M. Rangaswamy, *Trans. A.M.S.*, to appear)

$L_K(E)$ is primitive \Leftrightarrow

Primitive Leavitt path algebras

Can we describe the (left) primitive Leavitt path algebras?

Note: Since $L_K(E) \cong L_K(E)^{op}$, left primitivity and right primitivity coincide. So we can just say “primitive” Leavitt path algebra.

Theorem. (A-, Jason Bell, K.M. Rangaswamy, *Trans. A.M.S.*, to appear)

$L_K(E)$ is primitive \Leftrightarrow

1 $L_K(E)$ is prime,

Primitive Leavitt path algebras

Can we describe the (left) primitive Leavitt path algebras?

Note: Since $L_K(E) \cong L_K(E)^{op}$, left primitivity and right primitivity coincide. So we can just say “primitive” Leavitt path algebra.

Theorem. (A-, Jason Bell, K.M. Rangaswamy, *Trans. A.M.S.*, to appear)

$L_K(E)$ is primitive \Leftrightarrow

- 1 $L_K(E)$ is prime,
- 2 every cycle in E has an exit (Condition (L)),

Primitive Leavitt path algebras

Can we describe the (left) primitive Leavitt path algebras?

Note: Since $L_K(E) \cong L_K(E)^{op}$, left primitivity and right primitivity coincide. So we can just say “primitive” Leavitt path algebra.

Theorem. (A-, Jason Bell, K.M. Rangaswamy, *Trans. A.M.S.*, to appear)

$L_K(E)$ is primitive \Leftrightarrow

- 1 $L_K(E)$ is prime,
- 2 every cycle in E has an exit (Condition (L)), and
- 3 E has the Countable Separation Property.

$L_K(E)$ primitive $\Leftrightarrow E$ has (MT3), (L), and CSP

Strategy of Proof:

1. A unital ring R is left primitive if and only if there is a left ideal $N \neq R$ of R such that for every nonzero two-sided ideal I of R , $N + I = R$.

$L_K(E)$ primitive $\Leftrightarrow E$ has (MT3), (L), and CSP

Strategy of Proof:

1. A unital ring R is left primitive if and only if there is a left ideal $N \neq R$ of R such that for every nonzero two-sided ideal I of R , $N + I = R$.

Idea: (\Leftarrow) Embed N in a maximal left ideal T (this is OK since R is unital). So ${}_R R/T$ is simple.

Then $\text{Ann}_R(R/T) \subseteq T$ (noted previously). Thus $N + \text{Ann}_R(R/T) \subseteq T$. If to the contrary $\text{Ann}_R(R/T) \neq \{0\}$, the hypotheses would yield $N + \text{Ann}_R(R/T) = R$, impossible.

(\Rightarrow) If M is the supposed simple having $\text{Ann}_R(M) = \{0\}$, write $M \cong R/T$ for some maximal left ideal T . (In particular $T \neq R$.) So if $I \neq \{0\}$ then $I \cdot R/T = R/T$, so that $I + T = R$.

$L_K(E)$ primitive $\Leftrightarrow E$ has (MT3), (L), and CSP

2. Embed a prime $L_K(E)$ in a unital algebra $L_K(E)_1$ in the usual way; primitivity is preserved.

$L_K(E)$ primitive $\Leftrightarrow E$ has (MT3), (L), and CSP

2. Embed a prime $L_K(E)$ in a unital algebra $L_K(E)_1$ in the usual way; primitivity is preserved.
3. Show that CSP allows us to build a left ideal in $L_K(E)_1$ with the desired properties.

$L_K(E)$ primitive $\Leftrightarrow E$ has (MT3), (L), and CSP

2. Embed a prime $L_K(E)$ in a unital algebra $L_K(E)_1$ in the usual way; primitivity is preserved.
3. Show that CSP allows us to build a left ideal in $L_K(E)_1$ with the desired properties.
4. Then show that the lack of the CSP implies that no such left ideal can exist in $L_K(E)_1$.

$L_K(E)$ primitive $\Leftrightarrow E$ has (MT3), (L), and CSP

2. Embed a prime $L_K(E)$ in a unital algebra $L_K(E)_1$ in the usual way; primitivity is preserved.
3. Show that CSP allows us to build a left ideal in $L_K(E)_1$ with the desired properties.
4. Then show that the lack of the CSP implies that no such left ideal can exist in $L_K(E)_1$.

We will use:

“Reduction Theorem”. If E has Condition (L) then every nonzero two-sided ideal of E contains a vertex.

$L_K(E)$ primitive $\Leftrightarrow E$ has (MT3), (L), and CSP

(\Leftarrow). Suppose E downward directed, E has Condition (L), and E has CSP.

Suffices to establish primitivity of $L_K(E)_1$. Let T denote a set of vertices w/resp. to which E has CSP.

T is countable: label the elements $T = \{v_1, v_2, \dots\}$.

$L_K(E)$ primitive $\Leftrightarrow E$ has (MT3), (L), and CSP

Inductively define a sequence $\lambda_1, \lambda_2, \dots$ of paths in E for which, for each $i \in \mathbb{N}$,

- 1 λ_i is an initial subpath of λ_j whenever $i \leq j$, and
- 2 $v_i \geq r(\lambda_i)$.

Define $\lambda_1 = v_1$.

Suppose $\lambda_1, \dots, \lambda_n$ have the indicated properties. By downward directedness, there is u_{n+1} in E^0 for which $r(\lambda_n) \geq u_{n+1}$ and $v_{n+1} \geq u_{n+1}$. Let $p_{n+1} : r(\lambda_n) \rightsquigarrow u_{n+1}$.

Define $\lambda_{n+1} = \lambda_n p_{n+1}$. □

$L_K(E)$ primitive $\Leftrightarrow E$ has (MT3), (L), and CSP

Since λ_j is an initial subpath of λ_t for all $i \leq t$, we get that

$$\lambda_i \lambda_i^* \lambda_t \lambda_t^* = \lambda_t \lambda_t^* \quad \text{for each pair of positive integers } i \leq t.$$

In particular $(1 - \lambda_i \lambda_i^*) \lambda_t \lambda_t^* = 0$ for $i \leq t$.

$L_K(E)$ primitive $\Leftrightarrow E$ has (MT3), (L), and CSP

Since λ_i is an initial subpath of λ_t for all $i \leq t$, we get that

$$\lambda_i \lambda_i^* \lambda_t \lambda_t^* = \lambda_t \lambda_t^* \quad \text{for each pair of positive integers } i \leq t.$$

In particular $(1 - \lambda_i \lambda_i^*) \lambda_t \lambda_t^* = 0$ for $i \leq t$.

Define $N = \sum_{i=1}^{\infty} L_K(E)_1 (1 - \lambda_i \lambda_i^*)$.

$N \neq L_K(E)_1$: otherwise, $1 = \sum_{i=1}^t r_i (1 - \lambda_i \lambda_i^*)$ for some $r_i \in L_K(E)_1$, but then

$$0 \neq 1 \cdot \lambda_t \lambda_t^* = \left(\sum_{i=1}^t r_i (1 - \lambda_i \lambda_i^*) \right) \cdot \lambda_t \lambda_t^* = 0.$$

$L_K(E)$ primitive $\Leftrightarrow E$ has (MT3), (L), and CSP

Claim: Every nonzero two-sided ideal I of $L_K(E)_1$ contains some $\lambda_n \lambda_n^*$.

$L_K(E)$ primitive $\Leftrightarrow E$ has (MT3), (L), and CSP

Claim: Every nonzero two-sided ideal I of $L_K(E)_1$ contains some $\lambda_n \lambda_n^*$.

Idea: E is downward directed, so $L_K(E)$, and therefore $L_K(E)_1$, is prime. Since $L_K(E)$ embeds in $L_K(E)_1$ as a two-sided ideal, we get $I \cap L_K(E)$ is a nonzero two-sided ideal of $L_K(E)$. So Condition (L) gives that I contains some vertex w .

$L_K(E)$ primitive $\Leftrightarrow E$ has (MT3), (L), and CSP

Claim: Every nonzero two-sided ideal I of $L_K(E)_1$ contains some $\lambda_n \lambda_n^*$.

Idea: E is downward directed, so $L_K(E)$, and therefore $L_K(E)_1$, is prime. Since $L_K(E)$ embeds in $L_K(E)_1$ as a two-sided ideal, we get $I \cap L_K(E)$ is a nonzero two-sided ideal of $L_K(E)$. So Condition (L) gives that I contains some vertex w .

Then $w \geq v_n$ for some n by CSP. But $v_n \geq r(\lambda_n)$ by construction, so $w \geq r(\lambda_n)$. So $w \in I$ gives $r(\lambda_n) \in I$, so $\lambda_n \lambda_n^* \in I$.

Now we're done. Show $N + I = L_K(E)_1$ for every nonzero two-sided ideal I of $L_K(E)_1$. But $1 - \lambda_n \lambda_n^* \in N$ (all $n \in \mathbb{N}$) and $\lambda_n \lambda_n^* \in I$ (some $n \in \mathbb{N}$) gives $1 \in N + I$.

$L_K(E)$ primitive $\Rightarrow E$ has (MT3), (L), and CSP

For the converse:

$L_K(E)$ primitive $\Rightarrow E$ has (MT3), (L), and CSP

For the converse:

1) E not downward directed $\Rightarrow L_K(E)$ not prime $\Rightarrow L_K(E)$ not primitive.

$L_K(E)$ primitive $\Rightarrow E$ has (MT3), (L), and CSP

For the converse:

1) E not downward directed $\Rightarrow L_K(E)$ not prime $\Rightarrow L_K(E)$ not primitive.

2) General ring theory result: If R is primitive and $f = f^2$ is nonzero then fRf is primitive.

If E contains a cycle c (based at v) without exit then $vL_K(E)v \cong K[x, x^{-1}]$, which is not primitive, so $L_K(E)$ is not primitive.

$L_K(E)$ primitive $\Rightarrow E$ has (MT3), (L), and CSP

3) (The hard part.) Show if E does not have CSP then $L_K(E)$ is not primitive.

$L_K(E)$ primitive $\Rightarrow E$ has (MT3), (L), and CSP

3) (The hard part.) Show if E does not have CSP then $L_K(E)$ is not primitive.

Lemma. Let N be a left ideal of a unital ring A . If there exist $x, y \in A$ such that $1 + x \in N$, $1 + y \in N$, and $xy = 0$, then $N = A$.

Proof: Since $1 + y \in N$ then $x(1 + y) = x + xy = x \in N$, so that

$$1 = (1 + x) - x \in N.$$

$L_K(E)$ primitive $\Rightarrow E$ has (MT3), (L), and CSP

We show that if E does not have CSP, then there does NOT exist a left ideal $N \neq L_K(E)_1$ for which $N + I = L_K(E)_1$ for all two-sided ideals I of $L_K(E)_1$.

To do this: assume N is such an ideal, show $N = L_K(E)_1$.

$L_K(E)$ primitive $\Rightarrow E$ has (MT3), (L), and CSP

We show that if E does not have CSP, then there does NOT exist a left ideal $N \neq L_K(E)_1$ for which $N + I = L_K(E)_1$ for all two-sided ideals I of $L_K(E)_1$.

To do this: assume N is such an ideal, show $N = L_K(E)_1$.

Strategy: If N has this property, then for each $v \in E^0$ we have $N + \langle v \rangle = L_K(E)_1$. So for each $v \in E^0$ there exists $y_v \in \langle v \rangle$, $n_v \in N$ for which $n_v + y_v = 1$. Let $x_v = -y_v$. This gives a set $\{x_v \mid v \in E^0\} \subseteq L_K(E)_1$ for which $1 + x_v \in N$ for all $v \in E^0$.

$L_K(E)$ primitive $\Rightarrow E$ has (MT3), (L), and CSP

Now show that the lack of CSP in E^0 forces the existence of a pair of vertices v, w for which $x_v \cdot x_w = 0$. (This is the technical part.)

Then use the Lemma.

$L_K(E)$ primitive $\Rightarrow E$ has (MT3), (L), and CSP

Key pieces of the technical part:

- 1 Every element ℓ of $L_K(E)$ can be written as $\sum_{i=1}^n k_i \alpha_i \beta_i^*$ for some $n = n(\ell)$, and paths α_i, β_i . In particular, we can “cover” all elements of $L_K(E)$ by specifying n and lengths of paths. This is a countable covering of $L_K(E)$. (Not a partition.)

$L_K(E)$ primitive $\Rightarrow E$ has (MT3), (L), and CSP

Key pieces of the technical part:

- 1 Every element ℓ of $L_K(E)$ can be written as $\sum_{i=1}^n k_i \alpha_i \beta_i^*$ for some $n = n(\ell)$, and paths α_i, β_i . In particular, we can “cover” all elements of $L_K(E)$ by specifying n and lengths of paths. This is a countable covering of $L_K(E)$. (Not a partition.)
- 2 Collect up the x_v according to this covering. Since E does not have CSP, then some specific subset in the cover does not have CSP.

$L_K(E)$ primitive $\Rightarrow E$ has (MT3), (L), and CSP

Key pieces of the technical part:

- 1 Every element ℓ of $L_K(E)$ can be written as $\sum_{i=1}^n k_i \alpha_i \beta_i^*$ for some $n = n(\ell)$, and paths α_i, β_i . In particular, we can “cover” all elements of $L_K(E)$ by specifying n and lengths of paths. This is a countable covering of $L_K(E)$. (Not a partition.)
- 2 Collect up the x_v according to this covering. Since E does not have CSP, then some specific subset in the cover does not have CSP.
- 3 Show that, in this specific subset Z , there exists $v \in Z$ for which the set

$$\{w \in Z \mid x_v x_w = 0\}$$

does not have CSP. In particular, this set is nonempty. Pick such v and w . Then we are done by the Lemma. □



von Neumann regular rings

Definition: R is *von Neumann regular* (or just *regular*) in case

$$\forall a \in R \exists x \in R \text{ with } a = axa.$$

(R is not required to be unital.)

von Neumann regular rings

Definition: R is *von Neumann regular* (or just *regular*) in case

$$\forall a \in R \exists x \in R \text{ with } a = axa.$$

(R is not required to be unital.)

Examples:

- 1 Division rings
- 2 Direct sums of matrix rings over division rings
- 3 Direct limits of von Neumann regular rings

R is regular $\Leftrightarrow R_1$ is regular.

Kaplansky's Question

“Kaplansky's Question”:

I. KAPLANSKY, *Algebraic and analytic aspects of operator algebras*, AMS, 1970.

Is every regular prime algebra primitive?

Kaplansky's Question

“Kaplansky's Question”:

I. KAPLANSKY, *Algebraic and analytic aspects of operator algebras*, AMS, 1970.

Is every regular prime algebra primitive?

Answered in the negative (Domanov, 1977), a group-algebra example. (Clever, but very ad hoc.)

Kaplansky's Question

Theorem. (A-, K.M. Rangaswamy 2010)

$L_K(E)$ is von Neumann regular $\Leftrightarrow E$ is acyclic.

Kaplansky's Question

Theorem. (A-, K.M. Rangaswamy 2010)

$L_K(E)$ is von Neumann regular $\Leftrightarrow E$ is acyclic.

Idea of Proof: (\Leftarrow) If E contains a cycle c based at v , can show that $a = v + c$ has no “regular inverse”.

(\Rightarrow) Show that if E is acyclic then every element of $L_K(E)$ can be trapped in a subring of $L_K(E)$ which is isomorphic to a finite direct sum of finite matrix rings.

Application to Kaplansky's question

It's not hard to find acyclic graphs E for which $L_K(E)$ is prime but for which C.S.P. fails.

Example (mentioned previously): X uncountable, S the set of finite subsets of X . Define the graph E :

- vertices indexed by S , and
- edges induced by proper subset relationship.

Then for this graph E ,

- 1 $L_K(E)$ is regular (E is acyclic)
- 2 $L_K(E)$ is prime (E is downward directed)
- 3 $L_K(E)$ is not primitive (E does not have CSP).

Application to Kaplansky's question

Note: Embedding $L_K(E)$ in $L_K(E)_1$ in the usual way gives unital, regular, prime, not primitive algebras.

Remark: These examples are also “Cohn path algebras”.

Application to Kaplansky's question

A second construction of such graphs:

Let $\kappa > 0$ be any ordinal. Define E_κ as follows:

$$E_\kappa^0 = \{\alpha \mid \alpha < \kappa\}, \quad E_\kappa^1 = \{e_{\alpha,\beta} \mid \alpha, \beta < \kappa, \text{ and } \alpha < \beta\},$$

$s(e_{\alpha,\beta}) = \alpha$, and $r(e_{\alpha,\beta}) = \beta$ for each $e_{\alpha,\beta} \in E_\kappa^1$.

Application to Kaplansky's question

A second construction of such graphs:

Let $\kappa > 0$ be any ordinal. Define E_κ as follows:

$$E_\kappa^0 = \{\alpha \mid \alpha < \kappa\}, \quad E_\kappa^1 = \{e_{\alpha,\beta} \mid \alpha, \beta < \kappa, \text{ and } \alpha < \beta\},$$

$s(e_{\alpha,\beta}) = \alpha$, and $r(e_{\alpha,\beta}) = \beta$ for each $e_{\alpha,\beta} \in E_\kappa^1$.

Suppose κ has *uncountable cofinality*. Then $L_K(E_\kappa)$ is regular, prime, not primitive.

1 Primitive Leavitt path algebras

2 Primitive graph C^* -algebras

Prime graph C^* -algebras

For a ring R with a topology in which multiplication is continuous, then R is prime as a ring iff R is prime with respect to closed ideals. So for a C^* -algebra, primeness as a ring and primeness in the usual C^* sense mean the same thing.

Prime graph C^* -algebras

For a ring R with a topology in which multiplication is continuous, then R is prime as a ring iff R is prime with respect to closed ideals. So for a C^* -algebra, primeness as a ring and primeness in the usual C^* sense mean the same thing.

Proposition. Let E be any graph. Then $C^*(E)$ is prime if and only if

- 1 E is downward directed, and
- 2 E satisfies Condition (L).

Proof. This was established by Takeshi Katsura (2006), in the more general context of topological graphs.

$C^*(E)$ prime $\Leftrightarrow E$ has (MT3) and (L)

Idea of Proof:

Suppose E is downward directed and has (L).

If I and J are nonzero ideals in $C^*(E)$, then (L) with the Cuntz Krieger Uniqueness Theorem gives $u, v \in E^0$ such that $p_u \in I$ and $p_v \in J$.

Then downward directed gives $w \in E^0$ such that $u \geq w$ and $v \geq w$. So $p_w \in I$ and $p_w \in J$, so $0 \neq p_w = p_w^2 \in IJ$.

$C^*(E)$ prime $\Rightarrow E$ has (MT3) and (L)

Conversely: Suppose E does not satisfy (L). Then there exists a cycle $\alpha = e_1 \dots e_n$ in E without exits. If $H = \alpha^0$, then $I_H = I_{\overline{H}}$ is Morita equivalent to $C^*(\mathbb{T})$.

But this is impossible, since

- 1 any ideal of a prime C^* -algebra is itself prime as a C^* -algebra,
- 2 primeness is preserved under Morita equivalence, and
- 3 $C^*(\mathbb{T})$ is easily shown to not be prime.

So E satisfies Condition (L).

$C^*(E)$ prime $\Rightarrow E$ has (MT3) and (L)

Now show E is downward directed. Let $u, v \in E^0$. For $w \in E^0$

$$H(w) := \{x \in E^0 : w \geq x\}.$$

Let $\overline{H(w)}$ denote the saturated closure of $H(w)$.

For $u, v \in E^0$, the ideals $I_{H(u)} = I_{\overline{H(u)}}$ and $I_{H(v)} = I_{\overline{H(v)}}$ are each nonzero.

Since $C^*(E)$ is prime, $I_{\overline{H(u)}} \cap I_{\overline{H(v)}} \neq \{0\}$.

But $I_{\overline{H(u)} \cap \overline{H(v)}} = I_{\overline{H(u)}} \cap I_{\overline{H(v)}}$, so $\overline{H(u)} \cap \overline{H(v)} \neq \emptyset$, which gives that $H(u) \cap H(v) \neq \emptyset$.

Then $w \in H(u) \cap H(v)$ works.

Prime graph C^* -algebras

So the “answer” to the primeness question in the graph C^* -algebra setting differs from that of the Leavitt path algebra setting.

For example:

$$K[x, x^{-1}] = L(\bullet \curvearrowright) \text{ is prime,}$$

but

$$C^*(\mathbb{T}) = C^*(\bullet \curvearrowright) \text{ is not prime.}$$

Primitive C^* -algebras

Definition. The C^* -algebra A is *primitive* if there exists an irreducible faithful $*$ -representation of A .

Rephrased: A is primitive if there is an irreducible faithful representation $\pi : A \rightarrow B(\mathcal{H})$ as bounded operators on a Hilbert space \mathcal{H} . □

Primitive C^* -algebras

This will be useful:

Proposition: Suppose A is a C^* -algebra. Suppose there exists a modular left ideal $N \neq A$ of A such that $N + I = A$ for every nonzero closed two-sided ideal I of A . Then A is left primitive.

Primitive C^* -algebras

Idea of Proof. Suppose u is a modular element for N ; so
 $a - au \in N$ for all $a \in A$.

Standard: $u \notin N$ (otherwise $N = A$).

Standard: N embeds in a maximal (necessarily modular) left ideal T of A .

Standard: T is closed.

Primitive C^* -algebras

Idea of Proof. Suppose u is a modular element for N ; so $a - au \in N$ for all $a \in A$.

Standard: $u \notin N$ (otherwise $N = A$).

Standard: N embeds in a maximal (necessarily modular) left ideal T of A .

Standard: T is closed.

Since T is maximal, A/T is irreducible. Using closure of T and approximate identities for elements of A , standard to show that $\text{Ann}_A(A/T) \subseteq T$.

Now argue as in the unital ring case.

Primitive C^* -algebras

Lemma (well-known): Any primitive C^* -algebra is prime.

Primitive C^* -algebras

Lemma (well-known): Any primitive C^* -algebra is prime.

Proof. Let $\pi : A \rightarrow B(\mathcal{H})$ be the supposed irreducible faithful representation of the C^* -algebra A , and let I, J be (closed) two-sided ideals of A . Suppose $IJ = \{0\}$; we show that either $I = \{0\}$ or $J = \{0\}$. If $J \neq \{0\}$ then the faithfulness of π gives $\pi(J)\mathcal{H} \neq \{0\}$. But $\pi(J)\mathcal{H}$ is then a nonzero closed subrepresentation of the irreducible representation π , so $\pi(J)\mathcal{H} = \mathcal{H}$. Then $\{0\} = IJ$ gives $\{0\} = \pi(IJ)\mathcal{H} = \pi(I)\pi(J)\mathcal{H} = \pi(I)\mathcal{H}$, so that, again invoking the faithfulness of π , we get $I = \{0\}$. \square

Primitive C^* -algebras

Theorem (Dixmier, 1960) Every prime separable C^* -algebra is primitive.

Remark: It's an existence proof; the faithful irreducible representation is not explicitly constructed.

Primitive C^* -algebras

Theorem (Dixmier, 1960) Every prime separable C^* -algebra is primitive.

Remark: It's an existence proof; the faithful irreducible representation is not explicitly constructed.

Consequence: Suppose E is a graph for which $C^*(E)$ is separable. (So in particular E^0 is countable.)

Then $C^*(E)$ is primitive if and only if

Primitive C^* -algebras

Theorem (Dixmier, 1960) Every prime separable C^* -algebra is primitive.

Remark: It's an existence proof; the faithful irreducible representation is not explicitly constructed.

Consequence: Suppose E is a graph for which $C^*(E)$ is separable. (So in particular E^0 is countable.)

Then $C^*(E)$ is primitive if and only if

E is downward directed, and satisfies Condition (L).

Primitive C^* -algebras

Theorem (Dixmier, 1960) Every prime separable C^* -algebra is primitive.

Remark: It's an existence proof; the faithful irreducible representation is not explicitly constructed.

Consequence: Suppose E is a graph for which $C^*(E)$ is separable. (So in particular E^0 is countable.)

Then $C^*(E)$ is primitive if and only if

E is downward directed, and satisfies Condition (L).

... and, in this case, if and only if $L_K(E)$ is primitive.

Primitive graph C^* -algebras

Can we identify the primitive graph C^* -algebras for arbitrary graphs?

Note: “Primeness + Separability” of $C^*(E)$ is not the appropriate pairing of properties to achieve “Primitivity” in general.

For example $C^*(E)$ is primitive for E the graph with one vertex and uncountably many loops, but $C^*(E)$ is not separable.

Primitive graph C^* -algebras

Theorem. (A-, Mark Tomforde, in preparation)

Let E be any graph. Then $C^*(E)$ is primitive if and only if ...

- 1 E is downward directed,
- 2 E satisfies Condition (L), and

Primitive graph C^* -algebras

Theorem. (A-, Mark Tomforde, in preparation)

Let E be any graph. Then $C^*(E)$ is primitive if and only if ...

- 1 E is downward directed,
- 2 E satisfies Condition (L), and
- 3 E satisfies the Countable Separation Property.

Primitive graph C^* -algebras

Theorem. (A-, Mark Tomforde, in preparation)

Let E be any graph. Then $C^*(E)$ is primitive if and only if ...

- 1 E is downward directed,
- 2 E satisfies Condition (L), and
- 3 E satisfies the Countable Separation Property.

... if and only if $C^*(E)$ is prime and E satisfies the Countable Separation Property.

$C^*(E)$ primitive $\iff E$ has (MT3), (L), and CSP

Proof of sufficiency. *A lot of this will look familiar.*

Let X be a set of vertices with respect to which E satisfies the Countable Separation Property. Label the elements of X as $\{v_1, v_2, \dots\}$. We know (previous proof) there is a sequence $\lambda_1, \lambda_2, \dots$ of paths in E having the following properties for each $i \in \mathbb{N}$:

- (i) $v_i \geq r(\lambda_i)$, and
- (ii) $\lambda_{i+1} = \lambda_i \mu_{i+1}$ for some path μ_{i+1} in E .

$C^*(E)$ primitive $\iff E$ has (MT3), (L), and CSP

Proof of sufficiency. *A lot of this will look familiar.*

Let X be a set of vertices with respect to which E satisfies the Countable Separation Property. Label the elements of X as $\{v_1, v_2, \dots\}$. We know (previous proof) there is a sequence $\lambda_1, \lambda_2, \dots$ of paths in E having the following properties for each $i \in \mathbb{N}$:

- (i) $v_i \geq r(\lambda_i)$, and
- (ii) $\lambda_{i+1} = \lambda_i \mu_{i+1}$ for some path μ_{i+1} in E .

Note: since by construction $\lambda_1 = v$, $S_{\lambda_1} S_{\lambda_1}^* = P_v$.

$C^*(E)$ primitive $\iff E$ has (MT3), (L), and CSP

By construction, for $i < t$ we have

$$S_{\lambda_i} S_{\lambda_i}^* S_{\lambda_t} S_{\lambda_t}^* = S_{\lambda_t} S_{\lambda_t}^* \quad \text{for each pair of positive integers } i \leq t.$$

Claim: Every nonzero (closed) two-sided ideal J of $C^*(E)$ contains $S_{\lambda_n} S_{\lambda_n}^*$ for some $n \in \mathbb{N}$.

Reason: By Condition (L), the Cuntz-Krieger Uniqueness Theorem applies to yield that J contains some vertex projection P_w .

By the CSP there exists $v_n \in X$ for which $w \geq v_n$. But $v_n \geq r(\lambda_n)$.

So there is a path μ in E for which $s(\mu) = w$ and $r(\mu) = r(\lambda_n)$. Since $P_w \in J$ we get $P_{r(\lambda_n)} \in J$, so $S_{\lambda_n} S_{\lambda_n}^* = S_{\lambda_n} P_{r(\lambda_n)} S_{\lambda_n}^* \in J$.

$C^*(E)$ primitive $\Leftrightarrow E$ has (MT3), (L), and CSP

Let A denote $C^*(E)$, and let v denote v_1 . Consider the left ideal L of A defined by:

$$L = \left\{ \sum_{i=1}^n (x_i - x_i S_{\lambda_i} S_{\lambda_i}^*) \mid x_i \in A, n \in \mathbb{N} \right\}.$$

L is modular (with $a - aP_v \in L$ for all $a \in A$).

$P_v \notin L$. (Same proof as for Leavitt path algebras:)

$C^*(E)$ primitive $\iff E$ has (MT3), (L), and CSP

We use previous Proposition; need only show that $I + L = A$ for any nonzero closed two-sided ideal I of A . But any such two-sided ideal contains $S_{\lambda_n} S_{\lambda_n}^*$ for some $n \in \mathbb{N}$, hence contains $a S_{\lambda_n} S_{\lambda_n}^*$ for all $a \in A$, but then

$$a = a S_{\lambda_n} S_{\lambda_n}^* + (a - a S_{\lambda_n} S_{\lambda_n}^*) \in I + L.$$



$C^*(E)$ primitive $\Rightarrow E$ has (MT3), (L), and CSP

Proof of Converse.

Show that if $A = C^*(E)$ is primitive, then E has Condition (L), is downward directed, and has CSP.

$C^*(E)$ primitive $\Rightarrow E$ has (MT3), (L), and CSP

Proof of Converse.

Show that if $A = C^*(E)$ is primitive, then E has Condition (L), is downward directed, and has CSP.

Since primitive implies prime we get that E satisfies Condition (L) and is downward directed.

$C^*(E)$ primitive $\Rightarrow E$ has (MT3), (L), and CSP

So suppose to the contrary that E does not satisfy the Countable Separation Property. We show that $C^*(E)$ admits no faithful irreducible representations.

We actually show more, that $C^*(E)$ admits no faithful *cyclic* representations. Suppose $\psi : A \rightarrow B(\mathcal{H})$ is a cyclic representation of A ; so there exists $\xi \in \mathcal{H}$ for which $\psi(A)\mathcal{H} = \overline{\psi(A)\xi}$.

$C^*(E)$ primitive $\Rightarrow E$ has (MT3), (L), and CSP

We will use this general result:

Lemma. Let ψ be a representation of a C^* -algebra B as bounded operators on a Hilbert space \mathcal{H} , and let $\xi \in \mathcal{H}$. Suppose $\{Q_i \mid i \in I\}$ is a set of nonzero mutually orthogonal projections in B for which, for each $i \in I$, $\psi(Q_i)\xi \neq 0$. Then I is at most countable.

Proof. Use the Pythagorean Theorem in B .

$C^*(E)$ primitive $\Rightarrow E$ has (MT3), (L), and CSP

This graph-theoretic definition will also be useful.

Let E be any graph. For $w \in E^0$, let

$$U(w) = \{v \in E^0 \mid v \geq w\}.$$

Observation: E does *not* satisfy the Countable Separation Property in case for every countable subset X of E^0 , there exists some vertex v in $E^0 \setminus \bigcup_{x \in X} U(x)$.

$C^*(E)$ primitive $\Rightarrow E$ has (MT3), (L), and CSP

For every integer $n \geq 0$ define

$$\Gamma_n = \{\mu \in \text{Path}(E) \mid \psi(S_\mu S_\mu^*)\xi \neq 0, \text{ and } |\mu| = n\}.$$

(We view paths of length 0 as vertices, and in this case interpret $S_\mu S_\mu^*$ as $P_{s(\mu)\cdot}$)

Because the paths in Γ_n are of fixed length, the set $\{S_\mu S_\mu^* \mid \mu \in \Gamma_n\}$ consists of nonzero orthogonal projections.

So by the Lemma, each Γ_n is at most countable.

$C^*(E)$ primitive $\Rightarrow E$ has (MT3), (L), and CSP

For every integer $n \geq 0$ define

$$\Omega_n = \{w \in E^0 \mid w \in \mu^0 \text{ for some } \mu \in \Gamma_n\}.$$

Since each Γ_n is countable, and any path in E contains finitely many vertices, we get that each Ω_n is countable.

For every integer $n \geq 0$ define

$$\Theta_n = \bigcup_{w \in \Omega_n} U(w), \quad \text{and} \quad \Theta = \bigcup_{n=0}^{\infty} \Theta_n.$$

$C^*(E)$ primitive $\Rightarrow E$ has (MT3), (L), and CSP

Since $\Theta = \bigcup_{n=0}^{\infty} (\bigcup_{w \in \Omega_n} U(w))$, and each Ω_n is countable, we have that Θ is the union of a countable number of sets of the form $U(w)$.

So by the hypothesis that E does not satisfy the Countable Separation Property, we conclude that there exists some $v \in E^0 \setminus \Theta$.

$C^*(E)$ primitive $\Rightarrow E$ has (MT3), (L), and CSP

Since $\Theta = \bigcup_{n=0}^{\infty} (\bigcup_{w \in \Omega_n} U(w))$, and each Ω_n is countable, we have that Θ is the union of a countable number of sets of the form $U(w)$.

So by the hypothesis that E does not satisfy the Countable Separation Property, we conclude that there exists some $v \in E^0 \setminus \Theta$.

But $v \in E^0 \setminus \Theta$ means that for every path γ having $s(\gamma) = v$, then every path ν for which $r(\gamma) \in \nu^0$ has $\psi(S_\nu S_\nu^*)\xi = 0$.

$C^*(E)$ primitive $\Rightarrow E$ has (MT3), (L), and CSP

Let J denote the (nonzero) closed two-sided ideal of $C^*(E)$ generated by P_ν . Let $H(\nu)$ denote the set $\{w \in E^0 \mid \nu \geq w\}$.

Consider the set

$$T = \text{span}_{\mathbb{C}}\{S_\mu S_\nu^* \mid \mu, \nu \in \text{Path}(E) \text{ with } r(\mu) = r(\nu) \in H(\nu)\}.$$

Then T is dense in J .

$C^*(E)$ primitive $\Rightarrow E$ has (MT3), (L), and CSP

Claim: $\psi(t)\xi = 0$ for all $t \in T$.

Reason: Suffices to show that $\psi(S_\mu S_\nu^*)\xi = 0$ for any $\mu, \nu \in \text{Path}(E)$ for which $r(\mu) = r(\nu) \in H(\nu)$. But by the above description of $E^0 \setminus \Theta$ we have $\psi(S_\nu S_\nu^*)\xi = 0$, so that

$$\psi(S_\mu S_\nu^*)\xi = \psi(S_\mu S_\nu^* S_\nu S_\nu^*)\xi = \psi(S_\mu S_\nu^*)\psi(S_\nu S_\nu^*)\xi = \psi(S_\mu S_\nu^*)0 = 0.$$

$C^*(E)$ primitive $\Rightarrow E$ has (MT3), (L), and CSP

So $\psi(\overline{T})\xi = 0$, so that $\psi(\overline{T})\xi = 0$, and thus $\psi(J)\xi = 0$, which gives $\overline{\psi(J)\xi} = 0$. But then

$$\begin{aligned}\psi(J)\mathcal{H} &= \psi(J \cdot A)\mathcal{H} = \psi(J)\psi(A)\mathcal{H} = \psi(J)\overline{\psi(A)\xi} \\ &\subseteq \overline{\psi(J \cdot A)\xi} = \overline{\psi(J)\xi} = 0,\end{aligned}$$

so that $J \subseteq \text{Ker}(\psi)$. Since J is nonzero, ψ is not faithful. \square

Primitive C^* -algebras

We actually have shown more.

Definition. Let π be a representation of a C^* -algebra A on a Hilbert space \mathcal{H} . We say π is *countably generated* in case there exists a countable subset $\{h_i \mid i \in \mathbb{N}\}$ of \mathcal{H} for which

$$\mathcal{H} = \overline{\text{span}}\{\pi(A)h_i \mid i \in \mathbb{N}\}.$$

Primitive C^* -algebras

We actually have shown more.

Definition. Let π be a representation of a C^* -algebra A on a Hilbert space \mathcal{H} . We say π is *countably generated* in case there exists a countable subset $\{h_i \mid i \in \mathbb{N}\}$ of \mathcal{H} for which

$$\mathcal{H} = \overline{\text{span}}\{\pi(A)h_i \mid i \in \mathbb{N}\}.$$

Proposition. If E does not have CSP, then $C^*(E)$ admits no countably generated faithful representations.

Primitive C^* -algebras

Proof. Same idea as above. Suppose $\{h_i \mid i \in \mathbb{N}\} \subseteq \mathcal{H}$ has $\mathcal{H} = \overline{\text{span}}\{\pi(A)h_i \mid i \in \mathbb{N}\}$. For $n \geq 0, i \in \mathbb{N}$ define

$$\Gamma_n = \{\mu \in \text{Path}(E) \mid \psi(S_\mu S_\mu^*)\xi_i \neq 0 \text{ for some } i, \text{ and } |\mu| = n\}.$$

Now argue as before.

Prime, non-primitive C^* -algebras

The theorem gives us a machine to build prime, non-primitive C^* -algebras.

Example: The graph E as considered previously. X an uncountable set, S the set of finite subsets of X . E is the graph with:

- 1 vertices indexed by S , and
- 2 edges induced by proper subset relationship.

Then E is downward directed, has Condition (L), and does not have CSP.

So $C^*(E)$ is a prime, non-primitive C^* -algebra.

Note that $C^*(E)$ is an AF algebra.

Prime, non-primitive C^* -algebras

Modify E by adding a loop at each vertex. Call the new graph E' .

Then E' is still downward directed, has Condition (L), and does not have CSP.

So $C^*(E')$ is a prime, non-primitive C^* -algebra.

Note $C^*(E)$ is not AF. Also, since E' does not have Condition (K), $C^*(E)$ does not have real rank 0.

Prime, non-primitive C^* -algebras

Modify E' by adding a second loop at each vertex. Call the new graph E'' .

Then E'' is downward directed, has Condition (L), and does not have CSP.

So $C^*(E'')$ is a prime, non-primitive C^* -algebra.

Note that $C^*(E'')$ also has Condition (K), so has real rank 0.

Summary

Theorem. For an arbitrary graph E , these are equivalent.

- 1 E is downward directed, has Condition (L), and satisfies the Countable Separation Property.
- 2 $L_K(E)$ is primitive for every field K .
- 3 $L_{\mathbb{C}}(E)$ is primitive.
- 4 $C^*(E)$ is primitive.

Summary

Theorem. For an arbitrary graph E , these are equivalent.

- 1 E is downward directed, has Condition (L), and satisfies the Countable Separation Property.
- 2 $L_K(E)$ is primitive for every field K .
- 3 $L_{\mathbb{C}}(E)$ is primitive.
- 4 $C^*(E)$ is primitive.

Moreover, using this result, we can easily construct infinite classes of:

- 1 prime, non-primitive, von Neumann regular algebras, and
- 2 prime, non-primitive C^* -algebras.

Summary

Questions?